



Managing Pastures - Readers' Note

This document is part of a larger publication. The remaining parts and full version of the publication can be found at:

<http://www.dpi.nsw.gov.au/agriculture/livestock/dairy-cattle/feed/publications/managing-pastures>

Updated versions of this document can also be found at the above web address.

This document is subject to the disclaimers and copyright of the full version from which it is extracted. These disclaimers and copyright statements are available in the appropriate document at the above web address.

References

1. Curtis, A., Edmondson, C. and Findsen, C. 1989. Pasture Management for Dairy Farmers. Agriculture Victoria, Melbourne.
2. Donaghy, D., Scott, J. and Fulkerson, W. 1997. Grazing frequency affects *Lolium perenne* and *Trifolium repens* survival in the subtropics. Australian Journal of Experimental Agriculture (in press).
3. Fulkerson, W. J. 1994. Effect of redefoliation on the regrowth of water-soluble carbohydrate content of *Lolium perenne*. Australian Journal of Agricultural Research 45, 1809–15.
4. Michell, P. and Franks, D. 1993. The effects of grazing duration on herbage intake, composition, quality and regrowth of temperate pastures. Report on Dairy Research and Extension in Tasmania. Department of Primary Industries, Hobart.
5. Rogers, G. L., Porter, R. H. D. and Robinson, I. 1982. Comparison of ryegrass and white clover for milk production. In: Dairy Production from Pasture, Proceedings of the Conference of New Zealand and Australian Societies of Animal Production, Hamilton, NZ.
6. National Research Council 1989. Nutrient requirements of dairy cattle, 6th ed. National Academic Press, Washington, DC.
7. Davison, T. M. and Elliott, R. 1993. Response of lactating cows to grain-based concentrates in northern Australia. Tropical Grasslands 27, 229–37.

Questions

Grazing kikuyu at the 4–4½-leaf stage optimises quality. What is the penalty in quantity by not letting it grow longer?

If there is a high residue of ryegrass after grazing (well in excess of 5cm) for whatever reason, is it better to regraze or to wait for a full regrowth cycle?

Many ryegrass plants are lost through sod-pulling. What can be done to reduce this?

Is there any benefit in using nitrogen fertiliser on white clover to increase dry matter yields and take advantage of the high-quality herbage?

Is there any advantage in harrowing pastures to prevent dung pads, especially when short grazing rotations are being used?

When feed is in short supply, what value does concentrate have relative to silage?