

# GOA004 trapping of feral goats

Prepared by Trudy Sharp & Glen Saunders, NSW Department of Primary Industries

## Background

Feral goats (*Capra hircus*) can have a significant impact on the environment and agricultural production and are a potential reservoir and vector of endemic and exotic diseases. Although often considered a pest, feral goats are also an important resource, harvested commercially, primarily for meat. Control methods include trapping, mustering, exclusion fencing, ground shooting and shooting from helicopters.

Trapping of feral goats is usually conducted by landholders and involves the use of self-mustering technology which was originally developed for the management of sheep and cattle in rangeland areas. Trapping involves the construction of goat proof fences around water points with a number of one-way gates or ramps. The gates/ramps allow goats to enter the trap and have access to water but prevent them leaving. Once trapped, the goats are usually sold for live export, to abattoirs for slaughter or less commonly, for domestication, which offsets the costs of capture. Where there is no market for them or where removal may be costly or impractical e.g. in conservation areas or remote areas without access to transportation, the goats are usually destroyed by shooting in the trap yard.

This standard operating procedure (SOP) is a guide only; it does not replace or override the legislation that applies in the relevant State or Territory jurisdiction. The SOP should only be used subject to the applicable legal requirements (including OH&S) operating in the relevant jurisdiction.

## Application

- Trapping should only be used in a strategic manner as part of a co-ordinated program designed to achieve sustained effective control.
- Trapping is mainly used in semi-arid and arid rangelands where there are no alternative watering points for goats.
- Although traps can be costly to establish, trapping is more cost effective than mustering and is also less stressful for the goats. Trapping is the preferred method when goats are at low densities.
- Trapping is most effective during dry periods when goats drink regularly and are congregating around water holes. It becomes less effective and sometimes impractical during periods of wet weather when water is plentiful and goats are dispersed.

- Trapping at water can have significant negative impacts on non-target species, especially macropods and emus.
- Maintenance of traps is time consuming. Therefore, it is only suitable to use traps in situations where the operator has time to check them on a regular basis.
- Traps can also be used as self mustering yards for domestic stock such as sheep and cattle.
- Operators should endeavour to keep stress on the goats to a minimum during capture and handling. Prolonged stress not only has a negative impact on an animal's welfare but can also decrease carcass and meat quality.
- Shooting of goats should only be performed by skilled operators who have the necessary experience with firearms and who hold the appropriate licences and accreditation. Storage and transportation of firearms and ammunition must comply with relevant legislation requirements.

## Animal Welfare Considerations

### Impact on target animals

- Capture and handling increase stress in feral goats as they are not used to confinement or close contact with humans. Consequently, these procedures can result in mismothering, feeding disruption, social disruption, heat stress and also abortion in heavily pregnant females. Metabolic, nutritional and parasitic diseases and also changes in environmental conditions are common causes of mortality and morbidity in confined feral goats.
- Mixing unfamiliar animals may result in fighting, stress and/or injury. There should be sufficient yards to avoid mixing different groups of stock.
- Traps should be large enough to avoid overcrowding and allow all goats to access the water point.
- Feral goats should be handled quietly without force to avoid panic and trampling.
- Traps should be constructed to include trees, other vegetation and logs (located away from the edge of the fence line) to provide goats with shade and shelter. Goats can suffer when exposed to extremes of heat and cold.
- To avoid heat stress, trapping and handling of goats should be carried out when conditions are cool or mild.
- Capture and handling should be avoided when females are kidding or have dependent young at foot. Kids that do not accompany their mother into the trap may be separated and die of starvation or if trapped can get trampled underfoot. Although feral goats have been observed to breed at all times of the year, there are periods when the majority of kidding occurs e.g. in south-west Queensland, kidding mainly occurs in May to June. Generally, spring is the time of year when there is a greater proportion of does in late pregnancy or with young kids at foot.
- To minimise the possibility of starvation and stress, all traps must be inspected at least once daily. More frequent checking may be necessary during extreme weather conditions.

- The supply of water should be checked daily and appropriate feed must be made available if captured goats are to be held more than 24 hours. Account must be taken of their possible unwillingness to drink and eat from troughs. Animals being held for any length of time must be checked daily for ill thrift and signs of injury and disease.
- The trap should be constructed in a way so as not to cause injury from loose wire, sharp edges or malfunctioning gates. The trap gates should be large enough to allow big animals and those with large horns to enter the trap. The trap yard should be large enough so that each goat has enough space to avoid social stress.
- Fencing off alternative watering points to force goats to water at the trapped points has welfare implications. Some animals may not leave their preferred water source and will die of thirst rather than move and search for another.
- Goats that are found severely injured inside the trap must be killed quickly and humanely with a rifle shot to the head.
- Electric prods must not be used to assist in the handling of feral goats.
- Only non-aggressive, trained sheep dogs should be used to assist in the handling of feral goats, if used at all.
- Heavily pregnant, very young or weak/sick/injured animals must not be transported. They must either be destroyed, proper veterinary assistance given or transported at a later date when they are more suitable for transportation.
- The loading, transport, unloading, holding and slaughter of feral goats must be undertaken with the minimum amount of stress, pain or suffering. Guidelines on these procedures can be found in relevant state agriculture guidelines e.g.

NSW Agriculture Agfact A7.1.12 *Pre-slaughter management of goats* (2003) and in the following relevant Model Codes of Practice for the Welfare of Animals:

- The Goat (1991)
- Air Transport of Livestock (1986)
- Rail Transport of Livestock (1983)
- Road Transport of Livestock (1983)
- Sea Transport of Livestock (1987)
- Livestock at Slaughtering Establishments (2001)
- Killing or Capture, Handling and Marketing of Feral Livestock Animals (draft)
- Guidelines on the export of live goats can be found in the Australian Livestock Export Standards and the Livestock Export Accreditation Program Handbook (both available from LiveCorp at <http://livecorp.com.au/index1.cfm>).

## Impact on non-target animals

- Traps that would exclude large numbers of native species from natural springs and waterholes should not be constructed.
- If a trap continually catches numerous non-target animals, it should be constructed at another site where it will have minimal effect on other species or alternatively another goat control method could be used.

- Goat traps can have a significant negative impact on native non-target species by inadvertently trapping them and also by excluding them from water sources. For example macropods are affected by the following:
  - Traps that are closed from dusk to dawn will exclude macropods from drinking at a time when they mostly seek water;
  - Macropods may be reluctant to enter the trap even when the exit gates are open but will often hang around the perimeter fence trying to get access to water rather than moving on to another water source;
  - Macropods that enter an activated trap to drink will become trapped. Trapped macropods will rush at fences and injure themselves in an attempt to escape. They can also get caught in the fence when attempting to go over, through or under it; and
  - Macropods are not easily herded, so injuries and stress can be caused when trying to release those trapped out through the exit gates.
- Methods that could be used to reduce the impact of inadvertent trapping on non-target animals are a combination of engineering and management strategies. These include:
  - A barrier can be used on the external mesh fence to prevent kangaroos from getting their hind legs caught if they attempt to jump over. Chicken wire, rubber belting or shade cloth placed on the top 20 cm of the mesh acts as both a physical and visual barrier. The fence should be no more than 1.2 metres high (preferably 90cm).
  - Small escape gates can be incorporated at intervals around the fence to allow macropods to escape under the fence.
  - A protected water source could be provided nearby that would allow access to wildlife species but not to stock and feral goats.
  - Traps could be activated only during the day when goats and stock tend to water. This will help to avoid capture of macropods as they tend to water at night.
  - Moving macropods out of a trap should be done during the coolest part of the day to prevent them from overheating. Females should be closely monitored to see if they drop their pouch young. Macropods are very susceptible to capture myopathy, therefore they should be moved gently and quietly out of the yard through the trap gate prior to performing any other work in the vicinity of the trap.
- Trapped native non-target animals and also livestock that are still watering at the trapping point will need to be drafted from the trapped goats on a daily basis.
- To reduce the risk of injury to livestock, it is preferable to plan trapping sessions for times when livestock are out of the paddock (e.g. during shearing, lambing/calving, spelling). Trapping should be avoided during lambing/calving as ewes and cows can become separated from their young when they enter the trap for a drink
  - Non-target animals caught in traps must be examined for injuries and signs of illness or distress and dealt with as follows:
    - Animals which are unharmed or have only received minimal injuries such as minor cuts or abrasions should be immediately released at the site of capture.
    - Animals which have more severe injuries or which are suffering from thermal stress should receive appropriate attention. An animal suffering

from thermal stress can initially be placed in a suitable quiet holding area which provides warmth or shade to allow recovery before release. Animals with treatable injuries that cannot be immediately released or those failing to recover from thermal stress should be presented to a veterinarian or a registered wildlife carer for treatment.

- Animals that have injuries which are untreatable or which would compromise their survival in the wild should be euthanased using a technique that is suitable for the species. For more information on euthanasia techniques refer to **GEN001 *Methods of euthanasia***.

## Health and Safety Considerations

- Care must be taken when handling goats as they may carry diseases such as Q fever and scabby mouth (orf) that can affect humans and other animals. Routinely wash hands after handling goats or carcasses.
- Operators working with goats and goat carcasses are at risk of contracting Q fever. They may become infected when they inhale droplets of urine, milk, faeces or birth products from infected animals. Infection may also occur from inhalation of aerosols created during slaughter of infected animals or dust from contaminated materials. Blood testing of personnel is recommended to assess previous exposure, followed by vaccination for susceptible individuals
- During construction of traps, operators should be wary of the risks of injury from lifting heavy items. Leather gloves and eye protection will help prevent injuries from wire, steel posts and hammers.
- Firearms are potentially hazardous. All people should stand well behind the shooter when animals are being shot. The line of fire must be chosen to prevent accidents or injury from stray bullets or ricochets.
- Firearm users must strictly observe all relevant safety guidelines relating to firearm ownership, possession and use.
- Firearms must be securely stored in a compartment that meets state legal requirements. Ammunition must be stored in a locked container separate from firearms.
- Adequate hearing protection should be worn by the shooter and others in the immediate vicinity of the shooter. Repeated exposure to firearm noise can cause irreversible hearing damage.
- Safety glasses are recommended to protect eyes from gases, metal fragments and other particles.

## Equipment Required

### Traps

- Several trap designs exist, differing mainly in the one-way entrance. The three most commonly used traps are:

#### *Jump-down traps*

- The entrance consists of an earth ramp sloping up to approximately one metre high that allows the goats to access the trap by jumping down into it. A heavy gauge wire or baulking bar is placed approximately 30 cm above the top of the ramp to prevent the goats from jumping back out of the trap.

- The width of the ramp depends on the number of goats in the area.
- A gate is placed next to the ramp and this is left open when the traps are not in use to encourage the goats to use the traps.
- Jump-down ramps are best suited to areas that are free of livestock. Cattle and sheep that are in poor condition, and also lambs may suffer injuries when jumping from the ramp into the trap.
- Timid and small animals may be reluctant to use the ramp

#### *Spear gate traps*

- The entrance consists of a v-shaped, four-barred gate with flexible spears. Goats have to squeeze through the spears to enter the yard to drink.
- Goats have to be trained to go through the gates by gradually closing the spears to get them used to squeezing through.
- Big billies and other goats with large horns may have difficulty squeezing through this type of gate.

#### *Swinging one-way gate traps*

- These gates allow the goats to push through one way into the yard but do not move in the opposite direction when they push to get out.
- Trap yards should be large enough to comfortably handle the work they are expected to do. The most appropriate size will depend on the size of the water point, number and type of livestock using the water point, whether livestock and feral animals will be in the yard together and whether the animals will be held in the trap yard or drafted into holding yards. Large trap sizes give the goats enough room to move away from people entering the trap, allow for effective handling and will also reduce the pressure on and therefore damage to the fences. An adequate size to handle a large number of goats would be 50 × 50 metres.
  - It is preferable to incorporate loading pens and holding yards in the trap design that allows for on-site animal handling.
  - The yard fencing must be strong enough to withstand the pressure of animals bumping in to it. The most effective and economic fencing material used is ringlock or hingelock mesh. The most commonly used fence configuration is of prefabricated 8/90/15 hingelock with plain wires top and bottom to tie the hinged panels together i.e. the fence is 90 cm high, has 8 horizontal wires and a gap of 15 cm between vertical wires. The fence can also be topped with one or two plain wires and a strip of shade cloth material to increase the height of the fence (to no greater than 1.2 metres high).
  - Self-mustering trap yards have been built as squares, triangles and rounded yards. Round yards provide advantages over the other designs as the round shape provides the largest trap for material used, there are no corners to accumulate animals and the rounded shape aids in the flow of animals in, and through, the yard.
  - Choice of trap design will depend upon habitat, material available and accessibility to site. Knowledge of other species that may be at risk from inappropriately designed traps should be used to identify the most suitable trap designs and usage.
  - Details of trap specifications and construction can be obtained from relevant State agriculture guidelines, for example:

- Connelly, P. Horrocks, D., Pahl, L. and Warmin, K. (2000). *Cost effective and multipurpose self-mustering enclosures for stock*. Department of Primary Industries, Queensland.
- Henzell, R. (1984). *Methods of controlling feral goats*. Factsheet. Agdex 573. Department of Agriculture, South Australia.
- Joshua, E. (2003). *Yard design for goats*. Agfact A7.7.2. NSW Agriculture.
- King, D. (1991). *Goat control methods*. Infonote no. 21/91. Agriculture Protection Board of Western Australia.
- Underwood, C. (2002). *Total Grazing Management Field Guide: Self-mustering systems for cattle, sheep and goats*. Department of Agriculture, Western Australia.

## Firearms and ammunition

- Smaller calibre rifles such as .22 magnum rimfire with hollow/soft point ammunition are adequate for euthanasia of goats at short range (< 5 metres). If shooting animals from a greater distance refer to GOA001 *Ground shooting of feral goats* for more detailed information.

## Procedures

### Selection of trap sites

- Construct the trap at a site where there are limited numbers of watering points that can be fenced off easily. The trap should be situated on animal trails coming into the water point so that the gates are encountered on the usual path to water making it more likely the target species will quickly accept and continue to use it.
- If possible, choose a site that is in a shady area with as much natural vegetation as possible.
- Monitor the use of other watering points so that they can be fenced off if necessary to force goats to use the trap yard.
- Strategic placement is essential to reduce impact on local native species.

### Setting the trap

- If goats are being removed from the property for live sale, suitable transport must be arranged and confirmed before trapping is commenced.
- Prior to setting the trap an adequate training period (around 3 weeks) is required so that the all animals can become familiar with watering inside the trap yard. This period should be extended if any animals are showing difficulties in adapting.
- Once the animals are used to watering at the trap, the exit gate/s is/are closed and trapping commences.
- The trap should be checked at least once daily to avoid stress to the goats and to remove any domestic stock or non-target animals. Once trapped, goats are usually drafted into separate holding yards with access to feed and water. It is preferable to activate the trap each morning and then check in the evening.
- If bucks are observed to be fighting they should be drafted into separate yards.

- When checking the trap always approach from the direction of the gate. This will prevent the goats being forced into the gate area of the trap where the fence is lowest and there is a chance they may escape.
- Traps can be left at permanent sites with the gates open and reactivated when further trapping is required.

## Holding goats in yards

- Captured goats should be allowed a minimum of 48 hours rest with adequate shelter, food and water before they are transported on journeys longer than 8 hours. During this time they must be assessed daily for signs of injury, disease, inappetence, illness or distress. Account must be taken of their possible unwillingness to drink and eat from troughs.
- Goats should not be held in the trap or holding yards for extended periods. If goats are being held for any length of time (no longer than 3 days) they should be drafted into a large holding paddock that contains adequate shelter, food and water.
- To minimise injury in yards, ideally, goats should be segregated into the following groups:
  - Does with kids at foot;
  - Heavily pregnant does and small or young goats; and
  - Bucks

## Shooting of goats

- It may be necessary to humanely destroy goats by shooting in the following situations:
  - When there is no market for the captured goats (including smaller animals that are of no commercial value);
  - If goats have sustained serious injury during capture or in the holding yards;
  - Dependant young that are separated from their mother;
  - If goats have a previous disease or condition that would prevent them from being transported, slaughtered or domesticated.
- Shooting must be conducted to cause sudden and painless death with minimum distress to the animal. Only head shots are acceptable.
- The shooter should approach the animals in a calm and quiet manner. To prevent unnecessary agitation of the trapped goats, other people should keep away from the area until shooting is completed.
- To maximise the impact of the shot and to minimise the risk of misdirection the range should be as short as possible.
- Never fire when the goat is moving its head. Be patient and wait until the goat is motionless before shooting. Accuracy is important to achieve a humane death. One shot should ensure instantaneous loss of consciousness and rapid death without resumption of consciousness.
- Shots must be aimed to destroy the major centres at the back of the brain near the spinal cord. This can be achieved by one of the following methods (*see diagrams in Appendix*):

### Head Shot

The horn structures on adult goats make the temporal (side-on) or rear head shots the preferred points of aim. Shots to the front of the head can be used

on kids, however this method is not recommended for mature goats as the brain is located well back in the skull.

*Temporal position (side view)*

The firearm is aimed from the side of the head so that the bullet enters the skull at a point midway between the eye and the base of the ear on the same side of the head. The bullet should be directed horizontally into the skull.

*Rear of the head*

The firearm should be aimed at the back of the head at a point between the base of the horns and directed towards the mouth.

- Death of shot animals should always be confirmed by observing at least 3 of the following:
  - Absence of rhythmic, respiratory movements;
  - Absence of eye protection reflex (corneal reflex) or 'blink';
  - A fixed, glazed expression in the eyes; and
  - Loss of colour in mucous membranes (become mottled and pale without refill after pressure is applied).

If death cannot be verified, a second shot to the head should be taken immediately.

- When large numbers of animals are to be killed in the trap yard, provisions should be made to dispose of carcasses in an appropriate manner i.e. by burying and/or burning. Numerous guidelines are available which describe disposal methods e.g. Burton, 1999; AUSVETPLAN Operational Procedures Manual: Disposal (1996); NSW EPA (2001) Guidelines for disposal of dead stock.

## Further Information

Contact the relevant Commonwealth, State or Territory government agency from the following list of websites:

Commonwealth	Department of Environment and Heritage <a href="http://www.deh.gov.au/">http://www.deh.gov.au/</a>
ACT	Environment ACT <a href="http://www.environment.act.gov.au/">http://www.environment.act.gov.au/</a>
NSW	NSW Department of Primary Industries <a href="http://www.dpi.nsw.gov.au">www.dpi.nsw.gov.au</a>
NT	Parks & Wildlife Commission <a href="http://www.nt.gov.au/ipe/pwcnt/">www.nt.gov.au/ipe/pwcnt/</a>
QLD	Department of Natural Resources and Mines <a href="http://www.nrm.qld.gov.au">www.nrm.qld.gov.au</a>
SA	Animal & Plant Control Commission <a href="http://sustainableresources.pir.sa.gov.au">http://sustainableresources.pir.sa.gov.au</a>
TAS	Department of Primary Industries, Water & Environment <a href="http://www.dpiwe.tas.gov.au">www.dpiwe.tas.gov.au</a>
VIC	Department of Primary Industries, Agriculture & Food <a href="http://www.dpi.vic.gov.au">www.dpi.vic.gov.au</a>
WA	Agriculture WA <a href="http://www.agric.wa.gov.au">www.agric.wa.gov.au</a>

## References

Agriculture and Resource Management Council of Australia and New Zealand (1996). AUSVETPLAN (Australian Veterinary Emergency Plan). Operational Procedures Manual: Disposal. Edition 2 version 2.0. Document available electronically from the Animal Health Australia website: <http://www.aahc.com.au/ausvetplan/disfnl2.pdf>

American Veterinary Medical Association (2001). 2000 Report of the AVMA Panel on Euthanasia. *Journal of the American Veterinary Medical Association* **218**, 669–696.

Anon. (2001). Feral goat (*Capra hircus*). NRM facts PA18. Land Protection, Department of Natural Resources and Mines, Queensland.

Anon. (1999). Threat abatement plan for competition and land degradation by feral goats. The Environment and Heritage through the national threat abatement component of the Natural Heritage Trust. Document available electronically from the Department of the Environment and Heritage website: <http://www.deh.gov.au/biodiversity/threatened/tap/goats/index.html#contents>

Bellchambers, K. (2004). Improving the development of effective and humane trapping systems as a control method for feral goats in Australia. Project no. 40469. Department of the Environment and Heritage through the national threat abatement component of the Natural Heritage Trust.

Burton, R. (1999). Humane destruction and disposal of stock. Agnote DAI-136. NSW Agriculture, Orange. Document available electronically from the NSW Agriculture website: <http://www.agric.nsw.gov.au/reader/aw-companion/dai136.htm>

Casburn, G., Hacker, R and Brill, T. (1999). Evaluation of cooperative feral goat harvesting/control techniques. Project 94/1. Vertebrate Pest Program. NSW Agriculture, Orange.

Connelly, P., Horrocks, D., Pahl, L. and Warman, K. (2000). Cost effective and multipurpose self-mustering enclosures for stock. Department of Primary Industries, Queensland.

Finn, J., Greenwood, P. and May, T. (2003). Pre-slaughter management of goats Agfact A7.1.12. NSW Agriculture.

Henzell, R. (1984). Methods of controlling feral goats. Factsheet. Agdex 573. Department of Agriculture, South Australia.

Jago, B. (1999). Feral goat (*Capra hircus*) in Queensland. Pest status Review Series – Land Protection. Department of Natural Resources and Mines, Queensland.

Joshua, E. (2003). Yard design for goats. Agfact A7.7.2. NSW Agriculture.

King, D. (1991). Goat control methods. Infonote no. 21/91. Agriculture Protection Board of Western Australia.

Longair, J. A., Finley, G. G., Laniel, M. A., MacKay, C., Mould, K., Olfert, E. D., Roswell H. and Preston, A. (1991). Guidelines for euthanasia of domestic animals by firearms. *Canadian Veterinary Journal* **32**: 724–726.

NSW Environment Protection Authority (EPA) (2001). Procedure for dead stock disposal. Document available electronically from the NSW EPA website: <http://www.epa.nsw.gov.au/mao/deadstockdisposal.htm>.

O'Flynn, M. (1992). Animal welfare considerations. In D. Freudenberger (ed.), Proceedings of the National Workshop on Feral Goat Management: Planning for Action, Dubbo, NSW. Bureau of Resource Sciences, Canberra. pp 39–55.

Parkes, J., Henzell, R. and Pickles, G. (1996) *Managing Vertebrate Pests: Feral Goats*. Bureau of Resource Sciences and Australian Nature Conservation Agency. Australian Government Publishing Service, Canberra.

Ramsay, B. J. (1994). *Commercial use of wild animals in Australia*. Bureau of Resource Sciences. Australian Government Publishing Service, Canberra.

Standing Committee on Agriculture, Animal Health Committee. (1991). Model Code of Practice for the Welfare of Animals: Feral Livestock animals – Destruction or Capture, Handling and Marketing. CSIRO, Australia.

Thompson, J. Riethmuller, J. Kelly, D, Boyd-Law, S. and Miller, E. (1999). Technical report on feral goat management in south-west Queensland. Report to the Bureau of Rural Sciences. Queensland Department of Natural Resources.

Underwood, C. (2002). Total Grazing Management Field Guide: Self-mustering systems for cattle, sheep and goats. Bulletin No. 4547. Department of Agriculture Western Australia.

# Appendix

## Recommended shot placements - Feral goat

Diagram 1




Diagram 2 - Side view (skeleton)




Diagram 3 - Head shot (rear)



**Note:** Head shots (temporal or frontal) should be used for shooting feral goats at short range (< 5 metres). See text for details.



**Natural Heritage Trust**

*Helping Communities Helping Australia*  
A Commonwealth Government Initiative



NSW DEPARTMENT OF  
**PRIMARY INDUSTRIES**

## Disclaimer

The views and opinions expressed in this publication are those of the authors and do not necessarily reflect those of the Commonwealth and New South Wales Governments or the Commonwealth Minister for the Environment and Heritage and the New South Wales Minister for Primary Industries respectively. While reasonable efforts have been made to ensure that the contents of this publication are factually correct, the Commonwealth and New South Wales do not accept responsibility for the accuracy or completeness of the contents, and shall not be liable for any loss or damage that may be occasioned directly or indirectly through the use of, or reliance on, the contents of this publication.