



## **Hunter and Central Coast Sustainable Aquaculture Strategy**

### **Readers' Note**

---

This document is part of a larger publication. The remaining parts and full version of the publication can be found at:

<http://www.dpi.nsw.gov.au/fisheries/aquaculture/publications/general-management-and-policy/hunter-and-central-coast-sustainable-aquaculture-strategy>

Updated versions of this document can also be found at the above web address.

This document is subject to the disclaimers and copyright of the full version from which it is extracted. These disclaimers and copyright statements are available in the appropriate document at the above web address.

# Species Selection

Hunter and Central Coast Sustainable  
Aquaculture Strategy  
Land Based Aquaculture

A NSW Government Initiative

# Hunter and Central Coast Sustainable Aquaculture Strategy

A NSW Government initiative of Department of Primary Industries, Department of State and Regional Development, Department of Environment and Conservation, Department of Lands, Department of Infrastructure, Planning and Natural Resources and NSW Premiers Department to encourage sustainable aquaculture in New South Wales.

## Species Selection

### Table of Contents

1.	Selecting the Species	37
2.	Translocation Issues	37
3.	Hatchery and Genetic Factors	44
	3.1 Quality of Hatchery Progeny	44
	3.2 Broodstock Management	44
	3.3 Acquisition and Maintenance of Broodstock	44
	3.4 Broodstock Genetic Management	45
4.	Market and Pricing Factors	46
	4.1 There must be a market	46
	4.2 Factors affecting market value and price	47
	4.3 Product specification and adding-value	50
5.	Temperature Factors in Selecting Species	51
6.	Feed Factors	51
7.	Disease factors	52
	Reference Sources	55

## 1. Selecting the Species

The species selected for cultivation will affect the selection of a site and the design of the facility as well as the viability of the aquaculture business (*see also Site Selection and Planning & Design sections*). More than one species may be cultivated at a facility. In designing the facility, a certain amount of flexibility should be built into the design and layout so that the species can be switched later to meet opportunities created by changing markets.

In selecting species, the Translocation Policy to protect NSW's native fish populations must be considered. The selection of the species will also affect the level of assessment and type of approvals required.

Factors in the selection of species include:

- translocation policy;
- genetic factors, availability of seed stock (reliability, quality, quantity, season) and documented performance of the species in the aquaculture system proposed;
- market factors and potential profitability;
- site specific attributes (eg market distance, scale required, flood liability, temperature and water requirements (quality/quantity));
- feed and other management factors issues;
- disease factors.

In some situations, "polyculture" (i.e. more than one species within the one pond or in ponds in sequence) could provide opportunities for maximising returns and minimising environmental risk. In many circumstances, these types of multi-tiered systems can lead to more economical use of water, feed and energy, for example mullet grown in the prawn recirculation ponds.

## 2. Translocation Issues

One of the potential risks of aquaculture is the inadvertent introduction of species into waterways beyond their natural range or to areas within their natural range that have genetic stocks or populations that are distinct from the aquaculture stock. This is called translocation. Translocation of non-indigenous species can be sanctioned in some catchments, for example as a result of trout stocking for recreational fishing. In other circumstances, it may occur accidentally or deliberately but illegally.

Translocation of aquatic organisms has a number of inherent risks for the receiving aquatic habitats as well as for endemic organisms including:

- Predation;
- Introduction of disease or parasites;
- Competition, directly or indirectly for food and space;
- Adverse impacts on habitat and the native aquatic ecology; and
- Hybridisation with native fish that can affect genetic integrity.

The National Translocation Policy Guidelines has been developed to meet the needs of Australia's aquaculture and aquarium industries for the translocation of species within jurisdictions and across jurisdictional boundaries. These guidelines set out a risk assessment process for considering translocation issues. All proposals must be assessed according to the Translocation Policy.

Department of Primary Industries is currently refining the NSW Translocation Policy in the context of the National Translocation Policy Guidelines. However, for the purposes of this AIDP the principles in Table 6 and 7 also apply.

A number of the freshwater species considered in the AIDP are not indigenous to the region or the state and have therefore been assessed as a high risk species for the Hunter and Central Coast Region of NSW. High security species will have stricter requirements in terms of site selection, design and operational parameters. (See *Site Selection, Planning and Design and Operating the Farm sections*).

**Table 6. Key Translocation Principles in the Coastal Region**

1. Non-endemic species to NSW are not permitted to be translocated into estuarine pond systems.
2. Non-endemic species to NSW must meet prescribed health testing protocols for stock to be translocated from interstate.
3. Non-endemic species to NSW with high security status such as barramundi are permitted only in tank systems. Imported stocks must be disease free and discharge water treatment and disposal must meet the relevant AIDP performance criteria.
4. Non-endemic species to the Region with a high security status such as Murray Cod, Trout Cod, Barcoo Grunter and Sleepy Cod are only permitted if site selection, design and operational components meet the relevant AIDP performance criteria. Murray Cod and Trout Cod are not permitted in the Clarence and Richmond Catchments but may be cultured in other catchments.
5. Other species non-endemic to the Region such as Silver, Golden and Macquarie Perch are permitted in freshwater pond systems that meet the relevant AIDP performance criteria.
6. The threatened species Eastern Freshwater Cod may be permitted for culture in the Clarence and Richmond Catchments if sourced from accredited hatcheries that comply with Threatened Species Protocols.
7. Threatened species may be permitted for culture in their natural Catchments provided they are sourced from accredited hatcheries that comply with Threatened Species Protocols.

The generic risks associated with the culture of a particular species have been assessed and listed in Table 7 (*Species Table*).

The categorisation of each species was determined according to an assessment of the risks associated with translocation of disease and pathogens, and the risks associated with the survival and establishment of the species should it escape an aquaculture facility. The species are categorised into either a High or Low category as shown in Column 2 and Column 3 of Table 7. These categorisations are applied to set standards for facility design in the Project Profile Analysis under the heading Stock Security.

The Species Table also applies risk management measures that are specific to each species. Culture methods and locations that are considered too risky are prohibited in the Species Table.

**Table 7. Species Table**

Species	Disease/ Pathogen Security Status	Likely chance of Survival & risk of Establishing feral or invasive populations following escapement	Permissible culture methods <sup>1</sup>				Specific Operational and Locational Constraints
			Tanks and Raceways	Ponds Below PMF in the eastern drainage OR below 1:100 year flood level in western drainage.	Ponds Above PMF in the eastern drainage OR above 1:100 year flood level in western drainage.	Flow Through	
<i>Any hybrid fish OR</i> Any species not otherwise listed in this table	Assessment must be done on a case by case basis according to the National Policy on the Translocation of Live Aquatic Organisms						
<b>Freshwater aquarium species</b>							
Exotic freshwater aquarium species listed on Schedule 6 of the EPBC Act.	High	High	Yes	No	Yes	No	
Flat - headed gudgeon (Philypnodon grandiceps)	Natives:Low within endemic area - high outside	Natives:Low within endemic area - high outside/ domesticated natives: High	Yes	Yes	Yes	No	
Climbing galaxias (Galaxias brevipinnis)	As above	As above	Yes	Yes	Yes	No	
Common jollytail (Galaxias maculatus)	As above	As above	Yes	Yes	Yes	No	
Dwarf galaxias (Gallaxiella pusilla)	As above	As above	Yes	Yes	Yes	No	
Empire gudgeon (Hypseleotris compressa)	As above	As above	Yes	Yes	Yes	No	
Firetail gudgeon (Hypseleotris galii)	As above	As above	Yes	Yes	Yes	No	
Goldfish (Carassius auratus)	High	High	Yes	No	Yes	No	Prohibited in Catchments free of Carp
Koi carp (Cyprinus carpio)	High	High	Yes	No	Yes	No	Prohibited in Catchments free of Carp
Cox's gudgeon (Gobiomorphus coxii)	Natives:Low within endemic area - high outside	Natives:Low within endemic area - high outside/ domesticated natives: High	Yes	Yes	Yes	No	
Purple spotted gudgeon (Mogurnda adspersa)	As above	As above	Yes	Yes	Yes	No	
Murray cray (Euastacus armatus)	As above	As above	Yes	Yes	Yes	No	
Pacific blue eye (Pseudomugil signifer)	As above	As above	Yes	Yes	Yes	No	
Rainbow fish (Melanotaenia sp.)	As above	As above	Yes	Yes	Yes	No	
Striped gudgeon (Gobiomorphus australis)	As above	As above	Yes	Yes	Yes	No	
Sydney crayfish (Euastacus spinifer)	As above	As above	Yes	Yes	Yes	No	
Western carp gudgeon (Hypseleotris klunzingeri)	As above	As above	Yes	Yes	Yes	No	
Bullrout (Notesthes robusta)	As above	As above	Yes	Yes	Yes	No	
Freshwater mussels	As above	As above	Yes	Yes	Yes	No	
River blackfish (Gadopsis marmoratus)	As above	As above	Yes	Yes	Yes	No	
<b>Marine aquarium species</b>							
Barramundi cod (Cromileptes altivelis)	Low in east high in West	Low	Yes	No	No	No	

<sup>1</sup> For any **culture methods** not otherwise listed in this table an Assessment must be done on a case by case basis according to the National Policy on the Translocation of Live Aquatic Organisms.

**Hunter and Central Coast Sustainable Aquaculture Strategy  
Species Selection**

Species	Disease/ Pathogen Security Status	Likely chance of Survival & risk of Establishing feral or invasive populations following escapement	Permissible culture methods <sup>1</sup>				Specific Operational and Locational Constraints
			Tanks and Raceways	Ponds Below PMF in the eastern drainage OR below 1:100 year flood level in western drainage.	Ponds Above PMF in the eastern drainage OR above 1:100 year flood level in western drainage.	Flow Through	
Seahorse ( <i>Hippocampus kuda</i> )	As above	As above	Yes	No	No	No	
Seahorse ( <i>Hippocampus procerus</i> )	As above	As above	Yes	No	No	No	
Seahorse ( <i>Hippocampus trimaculatus</i> )	As above	As above	Yes	No	No	No	
Seahorse ( <i>Hippocampus tristis</i> )	As above	As above	Yes	No	No	No	
Seahorse ( <i>Hippocampus whitei</i> )	As above	As above	Yes	No	No	No	
Wrasse ( <i>Labroides bicolor</i> )	As above	As above	Yes	No	No	No	
Wrasse ( <i>Labroides dimidiatus</i> )	As above	As above	Yes	No	No	No	
Wrasse ( <i>Labroides pectoratus</i> )	As above	As above	Yes	No	No	No	
<b>Species non endemic to NSW</b>							
Atlantic salmon ( <i>Salmo salar</i> )	High	Low within present distribution or High elsewhere	Yes	Yes	Yes	Yes	Consistent with the EHNV zoning policy
Brook trout ( <i>Salvelinus fontinalis</i> )	High	Low within present distribution or High elsewhere	Yes	Yes	Yes	Yes	Consistent with the EHNV zoning policy
Brown trout ( <i>Salmo trutta</i> )	High	Low within present distribution or High elsewhere	Yes	Yes	Yes	Yes	Consistent with the EHNV zoning policy
Carp ( <i>Cyprinus carpio</i> )	High eastern / low western	High	Yes	Yes	Yes	No	Prohibited in Catchments free of Carp
Rainbow trout ( <i>Oncorhynchus mykiss</i> )	High	Low within present distribution or High elsewhere	Yes	Yes	Yes	Yes	Consistent with the EHNV zoning policy.
Redfin ( <i>Perca fluviatilis</i> )	High	High	Yes	Yes	Yes	Yes for catchments where they are present	Consistent with the EHNV zoning policy, for EHNV sentinel only Prohibited in Catchments free of Redfin
Barramundi ( <i>Lates calcarifer</i> )	High	low	Yes	No	No	No	Farms are to be above the PMF in the eastern drainage or above 1:100 year flood level in western drainage. No discharge of any waters to natural water bodies permitted.
Marron ( <i>Cherax tenuimanus</i> )	high	High	Yes	No	Yes	No	Special Fencing requirements
Redclaw ( <i>Cherax quadricarinatus</i> )	High	High	Yes	No	Yes	No	Special Fencing requirements
Sleepy cod ( <i>Oxyeleotris lineolatus</i> )	High	High	Yes	No	No	No	
Brine shrimp ( <i>Artemia sp.</i> )	High	High	Yes	Yes	Yes	No	
<b>Freshwater species</b>							
Australian bass ( <i>Macquaria novemaculeata</i> )	High in Western drainage Low in Eastern Drainage	High in Western drainage Low in Eastern Drainage	Yes	Yes	Yes	No	Eastern drainage sites - Broodstock must match DPI stocking zones

Hunter and Central Coast Sustainable Aquaculture Strategy  
Species Selection

Species	Disease/ Pathogen Security Status	Likely chance of Survival & risk of Establishing feral or invasive populations following escapement	Permissible culture methods <sup>1</sup>				Specific Operational and Locational Constraints
			Tanks and Raceways	Ponds Below PMF in the eastern drainage OR below 1:100 year flood level in western drainage.	Ponds Above PMF in the eastern drainage OR above 1:100 year flood level in western drainage.	Flow Through	
Eastern freshwater cod (Maccullochella ikei)	Low in Richmond and Clarence Drainage or High elsewhere	Low in Richmond and Clarence Drainage or High elsewhere	Yes	Yes	Yes	No	Prohibited outside of Richmond and Clarence Catchments.
Eel tailed catfish - Eastern Form (Tandanus tandanus)	High in Western drainage Low in Eastern Drainage	High	Yes	Yes	Yes	no	Stock to be sourced from approved Genetic Broodfish
Eel-long finned (Anguilla reinhardtii)	High in Western drainage Low in Eastern Drainage	High in Western drainage Low in Eastern Drainage	Yes	Yes	Yes	No	Pond design criteria
Eel-short finned (Anguilla australis)	High in Western drainage Low in Eastern Drainage	High in Western drainage Low in Eastern Drainage	Yes	Yes	Yes	No	Pond design criteria
Freshwater mullet (Myxus petardi)	High in Western drainage Low in Eastern Drainage	High in Western drainage & South of Shoalhaven - Low in remaining Eastern Drainage	Yes	Yes	Yes	No	
Cusped crayfish (Cherax cuspidatus)	High outside natural range / low within natural range	Low North Coast / High otherwise	Yes	Yes	Yes	No	Special Fencing required outside natural range
Rotund crayfish (Cherax rotundus)	High outside natural range / low within natural range	High outside natural range / low within natural range	Yes	Yes	Yes	No	Special Fencing required outside natural range
Strong crayfish (Euastacus valentulus)	High outside natural range / low within natural range	High outside natural range / low within natural range	Yes	Yes	Yes	No	Special Fencing required outside natural range
Freshwater prawn (Macrobrachium sp.)	High outside natural range / low within natural range	High	Yes	Yes	Yes	Yes	Broodstock must be sourced from local catchment
Freshwater shrimp (Atyidae sp.)	High outside natural range / low within natural range	High	Yes	Yes	Yes	Yes	Broodstock must be sourced from local catchment
Mussels (freshwater) (Vesunio ambiguus)	High outside natural range / low within natural range	High outside natural range / low within natural range	Yes	Yes	Yes	Yes	Broodstock must be sourced from local catchment
Barcoo grunter (Scortum barcoo)	High	High	Yes	No	No	No	
Bony bream (Nematalosa erebi)	High in Eastern drainage Low in Western Drainage	High in Eastern drainage Low in Western Drainage	Yes	Yes	Yes	No	
Eel tailed catfish - western form (Tandanus tandanus)	High in Eastern drainage Low in Western Drainage	High	Yes	Yes	Yes	No	Stock to be sourced from approved Genetic Broodfish
Golden perch (Macquaria ambigua)	High in Eastern drainage Low in Western Drainage	high (high genetic variation)	Yes	Yes	Yes		
Murray cod (Maccullochella peelii)	Low	Low	Yes	No in Eastern Drainage	Yes	No	Prohibited in Richmond and Clarence Catchments.
Silver perch (Bidyanus bidyanus)	High	high	Yes	Yes	Yes	No	
Spangled perch (Leiopotherapon unicolor)	High in Eastern drainage Low in Western Drainage	High in Eastern drainage Low in Western Drainage	Yes	Yes	Yes	No	
Welchs grunter (Bidyanus welchi)	High	High	Yes	No	No	No	
Yabby (Cherax destructor)	High in Eastern drainage Low in Western Drainage	High in Eastern drainage Low in Western Drainage	Yes	Yes	Yes	No	Special Fencing requirements in East
<b>Marine / estuary species</b>							
Balmain bugs (Ibacus peronii)	High in Western drainage Low in Eastern Drainage	low	Yes	Yes	Yes	Yes	
Banana prawn (Fenneropenaeus merguensis)	As Above	As Above	Yes	Yes	Yes	Yes	

Hunter and Central Coast Sustainable Aquaculture Strategy  
Species Selection

Species	Disease/ Pathogen Security Status	Likely chance of Survival & risk of Establishing feral or invasive populations following escapement	Permissible culture methods <sup>1</sup>				Specific Operational and Locational Constraints
			Tanks and Raceways	Ponds Below PMF in the eastern drainage OR below 1:100 year flood level in western drainage.	Ponds Above PMF in the eastern drainage OR above 1:100 year flood level in western drainage.	Flow Through	
Banded coral shrimp ( <i>Stenopus hispidus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Beachworm ( <i>Australonuphis parateres</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Beachworm ( <i>Australonuphis teres</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Beachworm ( <i>Hirsutonuphis mariahirsuta</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Black tiger prawn ( <i>Penaeus monodon</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Blacklip abalone ( <i>Haliotis rubra</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Bloodworms ( <i>Marphysa sanguinea</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Blue mussel ( <i>Mytilus gallo provincialis</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Brown tiger prawn ( <i>Penaeus esculentus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Clam ( <i>Tapes dorsatus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Cobia ( <i>Rachycteron canadum</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Coral trout ( <i>Plectropomus leopardus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Dusky flathead ( <i>Platycephalus fuscus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Eastern king prawn ( <i>Melicertus plebejus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Eastern lobster ( <i>Jasus verreauxi</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Estuarine clam ( <i>Katelysia rhytiphora</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Estuarine clam ( <i>Tapes dorsatus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Estuary cod ( <i>Epinephelus coioides</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Estuary perch ( <i>Macquaria colonorum</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Fingermark bream ( <i>Lutjanus johni</i> )	As Above	As Above	Yes	No	No	No	
Flat (mud) oysters ( <i>Ostrea angasi</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Flowery cod ( <i>Epinephelus fuscoguttatus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Greasyback prawn ( <i>Metapenaeus bennettiae</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Greenback flounder ( <i>Rhombosolea tapirina</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Kingfish ( <i>Seriola lalandi</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Kuruma prawn ( <i>Marsupenaeus japonicus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Mahi mahi ( <i>Coryphaena hippurus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Mangrove Jack ( <i>Lutjanus argentimaculatus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	

Hunter and Central Coast Sustainable Aquaculture Strategy  
Species Selection

Species	Disease/ Pathogen Security Status	Likely chance of Survival & risk of Establishing feral or invasive populations following escapement	Permissible culture methods <sup>1</sup>				Specific Operational and Locational Constraints
			Tanks and Raceways	Ponds Below PMF in the eastern drainage OR below 1:100 year flood level in western drainage.	Ponds Above PMF in the eastern drainage OR above 1:100 year flood level in western drainage.	Flow Through	
Mud crab ( <i>Scylla serrata</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Mullet ( <i>Mugil cephalus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Mulloway ( <i>Argyrosomus japonicus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Offshore greasyback prawn ( <i>M. ensis</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Pacific oysters ( <i>Crassostrea gigas</i> )	As Above	As Above	Yes	Yes	Yes	No	Noxious species in all estuaries except Port Stephens.
Queensland groper ( <i>Epinephelus lanceolatus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Red emperor ( <i>Lutjanus sebae</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Sand whiting ( <i>Sillago ciliata</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
School prawn ( <i>Metapenaeus macleayi</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Silver bream ( <i>Acanthopagrus australis</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Silver trevally ( <i>Pseudocaranx dentex</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Snapper ( <i>Pagrus auratus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Southern bream ( <i>Acanthopagrus butcheri</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Sydney rock oysters ( <i>Saccostrea glomerata</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Tubeworms ( <i>Diopatra aciculata</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Tubeworms ( <i>Diopatra dentata</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Yellow eye mullet ( <i>Aldrichetta forsteri</i> )	As Above	As Above	Yes	Yes	Yes	Yes	
Small mouth nannygai ( <i>Lutjanus erythropterus</i> )	As Above	As Above	Yes	Yes	Yes	Yes	

### 3. Hatchery and Genetic Factors

#### 3.1 Quality of Hatchery Progeny

Hatchery-reared juveniles can be used by the producers; sold to other aquaculture enterprises or the aquarium industry; or used for stocking into farm dams or for a Department of Primary Industries approved stock enhancement program in a natural waterway. It is the hatchery operators responsibility to ensure that:

- the juveniles are healthy before sale (it is illegal to sell diseased fish);
- the stock is not tainted with exotic invertebrates and vertebrates.

It is the hatchery's responsibility to ensure that the unintentional escape of cultured stock is minimised to prevent potential contamination of the genetic diversity of any endemic stocks and possible environmental impacts. Decisions relating to species and hatchery management will affect the planning and design of a facility and its operation (See *Hatchery* section in *Planning and Design*).

#### 3.2 Broodstock Management

The efficient production by hatchery facilities of quality, healthy juveniles in sufficient quantities to meet the increasing demands is dependent on the availability of adequate numbers of quality broodstock. The appropriate number of broodstock is determined by the number of eggs needed to produce the required number of juveniles, with normal losses taken into account. Other factors include the lifecycle of species, when broodstock are retired, genetic lines of broodstock and whether demand for juveniles is seasonal or continuous.

The assessment of the quality of broodstock usually focuses on factors such as:

- broodstock reproduction rates;
- performance and survival of broodstock under hatchery conditions;
- genetic characteristics; and
- progeny characteristics.

In the future as translocation issues and stock enhancement of wild fisheries becomes more important, hatchery breeding programs will be called upon to meet an increasing demands from the aquaculture industry as well as from specific stock enhancement programs or other fishery programs with more emphasis being placed on broodstock genetics.

#### 3.3 Acquisition and Maintenance of Broodstock

Broodstock can be collected from the wild under the provisions of a permit, grown and maintained in a hatchery or purchased from a commercial supplier. Currently, broodstock of Australian Bass are mostly captured in the wild prior to spawning, whereas broodstock of Silver and Golden Perch, Murray Cod, Snapper and Mulloway are maintained at the hatchery or in farm dams.

The reliability of spawning and quality of eggs from wild-caught broodstock varies with species. Reliance on the capture of wild broodstock limits the availability of eggs to the natural spawning period each year. Wild-caught broodstock are often used only once for spawning and then returned to the wild. Maintenance of broodstock in tanks with controls over day length and temperature can provide more flexibility in the supply of eggs and broodstock. For example, for many

species, the period of spawning can be extended from several months to year round.

Proper care of broodstock is essential for good production of eggs, larvae and juveniles. Methods differ with species, but the hatchery operators must provide optimum conditions for maturation and spawning. Factors for consideration include pond management, water quality and temperature, disease control, food supply, broodstock age, density and sex ratios.

### **3.4 Broodstock Genetic Management**

The characteristics of fish as with domesticated plants and animals can be improved by implementing breeding programs. For example, long-term breeding programs for chickens have resulted in significant improvements in productivity.

In any hatchery breeding program, a comprehensive record keeping system is necessary to track and evaluate progress of the program and to ensure weaknesses are not being introduced as a result of inbreeding.

#### **Selective Breeding Program to Improve Stock**

Fish stocks may be improved by selective breeding, hybridisation and crossbreeding. For example, selective breeding of trout has resulted in improved strains in relation to growth rates, spawning time and the age of maturation. Broodstock may be selected on a range of criteria such as size, colour, shape, growth, feed conversion rates, time of spawning, age at maturity, reproductive capacity and past survival rates. These criteria may vary with different hatcheries.

#### **Selective Breeding Program to Avoid Inbreeding**

Selective breeding involves selected mating of fish of a single strain and species and results in a reduction in the genetic variability in the population. Broodstock managers must be aware of the potential for inbreeding and manage the breeding program to minimise the risks of inbreeding problems arising. Inbreeding can occur when mates selected from a population of hatchery broodstock are closely related. The proportion of genes that fish parents have in common determines the degree to which a particular fish has been inbred. Problems that can occur after only one generation of brother-sister mating include reduced growth rates, lower survival, lower food conversion rates, increased numbers of deformed fry and increased susceptibility to disease. A system for selective breeding of trout has been developed to reduce the likelihood of inbreeding. The method involves maintaining three or more distinct breeding lines in a rotational line-crossing system. The number of fish required for the program can be set at any level necessary to satisfy the egg production requirements of a particular hatchery. Groups of fish can be held in separate tanks or combined, provided fish are individually tagged.

#### **Hybridisation and Cross Breeding Programs**

Hybridisation between species of fish and crossbreeding between strains of the same species can result in significant increases in growth, better food conversion, increased disease resistance and tolerance to environmental stressors. Hybrid vigour is responsible for the stock improvements; however this may not be achieved in all crosses.

### **Special Program Management for Stock Enhancement**

If hatchery-reared progeny are destined for a stock enhancement program, rather than aquaculture facility, maintenance of genetic diversity is extremely important. Broodstock should be obtained from the wild and mated pairs should be randomly selected from the population.

## **4. Market and Pricing Factors**

### **4.1 There must be a market**

In addition to a species being biologically suitable for aquaculture production, it must have market acceptance. The aim of commercial aquaculture is to make a profit. The farm must produce fish that are marketable in sufficient quantities at acceptable prices. All too often, a decision is made to farm a species based on biological factors with little consideration given to its market acceptance. These factors must be taken into consideration in the preparation of the Business Plan (see *Business Plan section*).

#### **Domestic market**

The live fish market in Australia is still developing, particularly for native freshwater species. There is still significant potential for expansion. However, once this market niche that returns very attractive prices has been filled, alternative higher volume markets are likely to return lower prices. In some cases, these prices may not be sufficient to cover production costs. If a number of farmers independently decide to target the live fish and other boutique markets at the same time, this may saturate the market leading to price wars and lower returns to farmers. It is important that farmers network with each other to ensure that a steady and reliable supply of quality product is supplied to meet market specifications and requirements. This approach will assist in increasing demand and help to smooth out periods of glut and short supply, thereby leading to more reliable prices for the farmed products. In addition, industry groups are better placed to develop and promote markets for new aquaculture products than individual producers.

A most important segment in the domestic market is Asian restaurateurs and fish retailers. By contrast the non-Asian retail trade is relatively small and more price sensitive. It is considered that this segment is capable of substantial growth with market promotion to improve consumer awareness.

In the longer term, it is considered that significant growth in the aquaculture fish "fillets" market will occur to meet the demand of the Australian consumers preference in the home and restaurant trade for skinless boneless fillets.

#### **Export markets**

There is excellent potential to promote the *clean green* image of Australian aquaculture products. In addition, it is anticipated that there will be great potential for Australian producers to step in and replace some of the shortfall in supply that may arise overseas. Australia, and in particular NSW are fortunate to have avoided most of the problems experienced by other countries associated with waterway pollution and disease. This *clean green* image is increasingly important because of growing concerns associated with the spread of disease and pest species with importing of fish especially live fish. As a result the development and implementation of best practice in production and quality assurance programs to guarantee the quality of the product is increasingly important for the expansion of the export market.

## 4.2 Factors affecting market value and price

There are many factors likely to affect the price achieved in the market place ranging from global economic issues (eg. value of the A\$ and the buoyancy of the Asian economy) to supply and demand factors (eg. competition from other fish products, consumer preference, the season and fads in eating). The price gained for a product will also vary with the market sector and its geographic location. For example there can be significant difference in price achieved in the local market and the Sydney, Brisbane and Melbourne markets and between wholesalers or overseas distributors, retailers (super markets and fishmongers), restaurateurs and the take-away-food sector.

It is important that the farmer becomes well informed about options and the implications in relation to the costs in getting the product to a particular market and the likely differentials in returns. For example:

- Where is the product to be positioned in the market place – economy or luxury?
- What are competitors offering and what prices are they achieving?
- What is the price range of potential customers and for what volume?
- Is there a potential for repeat business?

The lowest acceptable price should be equal to the cost per kilogram to produce the product, taking into consideration both fixed and variable costs. The highest price may be what is achievable by selling the best of the stock into a niche market. The base price is generally somewhere between the two.

**Table 8. Indicative prices from selected fish sales 2002**  
(*Ruello & Assoc pers comm*)

Fish/product whole or gilled and gutted (GG)	Wholesale or auction price approximate \$/kg	Median Sydney Retail price approximate \$/kg
Golden perch GG wild	13.50	20.00
Barramundi whole farm	8.50	14.00
Snapper GG wild	10.00	16.00
Silver perch whole farm	8.80	13.00
Silver bream whole wild	8.50	13.50
Rainbow trout GG farm	6.50	11.50

### MARKET FACTORS

#### Prawns

The Australian aquaculture production of prawns is estimated at about 2,700 t/year. Much is exported. Farm-gate prices vary around an average of \$15/kg for black tiger prawns, and up to as much as \$70/kg for live kuruma prawns sold on the Tokyo markets.

#### Crayfish

The size of the market for the three freshwater crayfish species grown in NSW is unknown. Redclaw have a meat yield (meat to body weight ratio) of 22 to 25 per cent; the meat yield is comparable in marron but slightly lower in the yabby. Yabbies have been marketed in small quantities from the wild fishery for years (up to \$8/kg). Much higher prices (up to \$35/kg) have been obtained for all three species by supplying live crayfish of a particular size to restaurants. These prices are for very small quantities, however prices in the range of \$8 to \$15 can be expected as supply increases. Crayfish are likely to feature in the highly competitive "entree" market (along with such delicacies as prawns, oysters and smoked salmon), and this may make it difficult to permanently establish a market niche. The size of any export

market is impossible to evaluate until supply can be maintained. There is certainly a market for crayfish in Europe, as crayfish are regarded as a delicacy and local stocks have been largely wiped out by the "crayfish plague". However, the market in some European countries (for example, Sweden) is highly seasonal. Trial shipments have suggested that *Cherax* species would be accepted in the marketplace.

#### **Silver Perch**

Most of the silver perch currently produced has been targeted for the live fish trade in Sydney's Asian restaurants. The live fish market requires fish to be 600-800g in weight and have a good physical appearance (no marks or blemishes on skin, fins intact, good colour etc). Silver perch has been readily accepted into this market, with farmers receiving strong prices (\$9-10/kg live) at the farm gate. It is expected that this price will drop in time as more producers enter the industry. Some live product is sold at the Brisbane and Melbourne Fish Markets, with small amounts sold regionally. Fish not suitable for the live trade is usually sold whole chilled on the Sydney Fish Market auction floor with a small quantity sold in the Brisbane and Melbourne Fish Markets.

In a recent report by Ruello Associates Pty. Ltd entitled "*Silver Perch Market Assessment*", it was considered that the market for whole silver perch is near saturated, and is only capable of growth of about 20-30% over the next few years. The report concluded that the long-term future for fish farming lies with fillets. The Australian consumer's preference is overwhelmingly for a skinless boneless fillet rather than a fish with its head and bones.

Local markets for fillets appear to be relatively unfulfilled with the potential domestic markets not yet fully explored with some researchers predicting that silver perch industry was capable of supplying over 10,000 tonnes/annum. There may also be future potential to supplement some of the \$500 million worth of fish imported into Australia every year, predominantly in the form of cheap, processed/frozen, white flesh fillets eg. hoki and hake. Export markets for silver perch are still relatively unexplored.

Like most freshwater fish, silver perch has a tendency to develop a muddy flavour. This occurs as a result of fatty tissues in the fish absorbing compounds released by blue green algae in the culture ponds. The only way to remove the *off-flavour* is by purging in clean water for 3-21 days, depending on the extent of tainting. Purging the fish in clean water expels the compounds that cause the taste, and the fish is then ready for market. The presence of off-flavours in marketed product has the potential to impact the industry. In order to encourage quality assurance and to assist maintain product standards, it is recommended that all silver perch are purged to reduce the potential for off-flavoured fish entering the markets. It is essential that the industry as a whole implement best management quality assurance practices so a high quality product is consistently available to the consumer.

#### **Barramundi**

One advantage to barramundi farmers is that the species has been commercially fished for a number of years, and as such has a well-established "elite" position in the market place. There are a number of options available to producers, including the live fish trade, plate size whole (300-500g) trade, and fillet or larger whole fish (2kg) trade. The reputation of barramundi as well as its premium edible properties (white, firm, mild tasting) also provides a good marketing platform for new producers. barramundi (cultured and wild-caught) is sold in most major fish market or retail outlets, with aquaculture barramundi usually in the form of the whole plate size product. Live fish are also sold, targeted at the Asian restaurant trade. Farmers receive an average price of around \$9-10/kg at the farm gate for Barramundi, with higher prices for value-added products eg fillets. The average price used to be higher, but has been driven down by the number of new producers entering the industry, as well as cheap imports from places such as Vietnam. Due to the higher production costs associated with barramundi culture compared to overseas producers, there is probably only limited potential to create export markets for this species.

### **Murray Cod**

Murray Cod has generated much interest in the live fish trade, due to its appearance and premium quality flesh and taste. Murray Cod sold as a live product has been very well received, and is perceived by some chefs to be one of the best tasting freshwater fish in the world. This is reflected in the average price paid for live Murray Cod; around \$25-30/kg at the farm gate. There is also good potential for a gilled/gutted product fetching prices of \$18-22/kg regularly on the Sydney Fish Markets auction floor. While most of the product is sold domestically, there may be potential for export trade of cod when production is increased. Overseas traders have expressed interest in the product. Most live product is purchased by Asian restaurants, indicating its acceptance within the Asian community. Although it is expected that average prices for live product will drop as more producers come on line, the almost legendary status of the cod, its excellent flavour and excellent appearance should ensure its place in the top range of the market.

### **Snapper**

While marine fish are more usually grown in cages in estuaries, bays or the ocean, pond-rearing trials have also produced a high quality product. Snapper are relatively easy to market. The majority of the existing demand is for plate-size fish (400g), which are sold and served whole. Two ways to expand the market for Australian snapper have been identified. The first is to export live and sashimi-quality snapper to Japan based on technology developed in New Zealand to export live wild snapper. The second is to develop a local market for live and sashimi snapper.

### **Mulloway**

Marketing mulloway is a different proposition to snapper. Large mulloway (more than 2.5kg) currently fetch the best price. Wild-caught smaller mulloway (soopies) have poorer consumer acceptance, however informal taste tests of 1kg mulloway grown in seacages were very favourable, and pond rearing trials have also produced a high quality product. Little is known of the export potential for mulloway.

### **Eels**

More than 130,000 tonnes/yr of eels is produced worldwide, with the main producers being China, Japan and Taiwan. Over 70% of this product is produced for the Japanese 'Kabayaki' market. Kabayaki is a style of serving eels, where eels of around 150-200g are butterflied, placed on skewers, basted in a thick soy based sauce, and steamed or grilled. More than 90% of eels consumed in Japan are served this way, with eel being the most widely consumed freshwater fish in Japan. The Australian shortfin eel (*Anguilla australis*) is very similar in appearance to the species of eel favoured by the Japanese market (*A. japonica*). As such, the shortfin eel is well accepted in Japan and attracts similar prices to *A. japonica*, averaging around \$10-15/kg at the farm gate (live). There is good potential for Australian producers to export shortfin eels to Japan for this market. The long fin eel (*A. reinhardtii*) is different in appearance to the *A. japonica* species, and as such is not well accepted in Japan, leading to lower prices.

However, the long fin eel is similar to the species favoured in China (*Anguilla marmorata*), and as such there is some potential to export this species there. For eels, it is hard to evaluate the size of any potential markets for Australian producers. Certainly the export markets are present and active; however the very nature of eel farming (i.e. reliance on glass eels for seedstock, seasonal nature of glass eel collection, limited technical know-how etc.) can to some extent prohibit its development. Declining eels stocks overseas may work to the advantage of Australian producers. Furthermore, as Australia's multicultural population increases, there may be opportunity for a small domestic market for eels to develop.

### **Trout**

Trout have high meat yields and sell well. Fresh trout has become cheaper than other prime fish such as snapper, mulloway and barramundi. They command medium prices for quality fish (rainbow trout retails for approximately \$8 per kg fresh and \$10 smoked; with approximately \$4-6 at the Sydney Market). At present, a few large producers dominate the market and control the maximum price. Value-added products (e.g. smoked trout) demand higher prices. New marketing techniques (e.g. fillets and cutlets in supermarkets) are being developed.

### 4.3 Product specification and adding-value

Like other primary industries, if the aquaculture industry is to be sustainable, it must constantly adapt to changing preferences and expectations of consumers, and to changing market conditions. To do this, the industry needs to focus on meeting the market and where possible value-adding. The following outlines marketing options:

- Live fish, prawns or other species (to remain alive for 5 days),
- Chilled whole fish (shelf life for 1 week at 5°C) (eg. GU=Gutted, GH=Headed/Gutted, GG=Gilled/Gutted),
- Cooked prawns and yabbies - whole or peeled – frozen or chilled,
- Skinless and boneless chilled or frozen fillets (FI=Fillets) – chunks, steaks, fillets, nuggets etc.
- Smoked (hot or cold smoking), pickled or other processed products.

The size of the fish can also be a key factor in market returns and consideration should be given to the grading specifications at the Sydney Markets which may change from time to time.

**Table 9. Current Grading Specifications at the Sydney Markets**

<b>Aquaculture species</b>	<b>Extra small</b>	<b>Small</b>	<b>Medium</b>	<b>Large</b>	<b>Extra large</b>
Barramundi <i>gms</i>		< 450	450 - 600	600 - 800	800 +
Silver Perch <i>gms</i>		300 - 450	450 - 550	> 550	
Black Tiger Prawns (P.monodon) <i>Number/kg</i>	>66	56 – 66	45 - 56	22 - 44	<21

Market sectors exhibit a preference for certain sized fish – for example for silver perch (Ruello & Assoc):

- Large whole fresh or live fish for the Asian “shared plate” dining range typically from 500gms – 1kg and up to over 2kg at times. There is likely to be greater demand for this size at holiday periods.
- Chilled whole fish for non-Asian cuisine ranging from 400gms – 1kg.
- Chilled plate-sized for Asian and non-Asian cuisine sold mostly whole or some gutted ranging from 350 - 500gms.
- Fillets mostly for non-Asian cuisine – 100 – 200gms fillets from fish 300 – 800gms.

Packaging and presentation of products into the retailing market is an issue that requires more consideration. Bulk fish packaging is usually in clean styrene cases with a polyethylene liner bag in 10kg or 15 - 18kg lots. Cooked king prawn supply is a mix of plastic tubs (20 - 25kg) and styrene cases (10kg and 16 - 18kg type).

Positioning speciality products through the use of well-designed packaging can add value and increase returns. The key to success with value-added products is a good quality assurance program to ensure that consistently high quality products reach the consumer. All products should meet the National Food Standards (See *Operating the Farm* section and Australian Seafood Industry Quality Assurance Project by Seafood Services Australia [www.dpi.qld.gov.au/cft/ssaust](http://www.dpi.qld.gov.au/cft/ssaust)).

## 5. Temperature Factors in Selecting Species

Water temperature profoundly affects the growth rates and health of fish. Most species of finfish that are farmed successfully reach minimum market sizes (40-500gms) in 12 -18 months. Prawns will typically reach market size in 3-6 months. To enable maximum growth, the temperature range of the selected site should match with the optimum temperature growing regime for the selected species. Regions where lethal temperatures are reached, or even approached, are unsuitable for pond or raceway culture of the particular species. Consideration of the site/system design in terms of likely water temperature ranges/growing season is essential for the species under consideration. Below is a table of optimal temperatures for various species under consideration that should be considered during site selection (See *Site Selection* section).

**Table 10. Temperature range for breeding and grow-out**

Species	Ideal Temperature	
	Hatcheries	Grow-out facilities
Abalone	15 - 18°C	12 - 20°C
Prawn - Black Tiger	28 - 32°C	25 - 32°C
Prawns - Kuruma	25 - 30°C	20 - 28°C
Prawns - School	-	21 - 27°C
Crustacea - Redclaw	27 - 30°C	27 - 32°C
Crustacea - Yabbies	15 - 20°C	23 - 25°C
Barramundi	27 - 30°C	26 - 30°C
Eels	-	23 - 28°C
Kingfish	21 - 24°C	15 - 25°C
Mahe-Mahe	25 - 30°C	25 - 30°C
Mulloway	21 - 26°C	14 - 30°C
Murray Cod	19 - 21°C	23 - 26°C
Silver Perch	20 - 25°C	23 - 28°C
Snapper	21 - 24°C	17 - 30°C
Trout Brown	*6 - 10°C	4 - 19°C
Trout Rainbow	*9 - 14°C	10 - 22°C

*\*for spawning and egg production*

## 6. Feed Factors

Intensive fish culture is based on the use of medium to high stocking densities, prepared feeds (usually specially formulated dry pellets) and, with some species, fertilisation of pond water. Very high production rates can only be achieved with species that accept prepared feed. Not all species readily accept pellet rations (e.g. Australian bass and golden perch) and consequently there has been limited progress with intensive cultivation of these species.

Species vary in their optimal protein and nutrient intake. Generally, more carnivorous species such as snapper and kuruma prawns require a higher animal protein component, whereas species with lower protein requirements such as silver perch and abalone are often more amenable to diets with inclusion of plant protein sources. For aquaculture to continue to expand significantly, fish meal based diets must be replaced with other protein sources.

The food conversion ratio (FCR) is the ratio of dry weight of food, to the wet weight gain of fish. The lower the ratio, the more efficiently food has been converted to fish flesh. Feed costs often constitute 40-55% of total production costs, so it is essential to use species that convert food efficiently. Modern diets enable food conversion ratios of 2:1 or better in freshwater species such as trout. Species that have high meat to total body weight ratio are desirable because of their more efficient conversion of feed into edible flesh. This is particularly important if the end product is to be processed.

## 7. Disease factors

Although all species are susceptible to disease under culture conditions, the interactions that cause disease outbreaks relate to three components referred to as the epidemiological triad namely:

- the disease agent;
- the host; and
- environmental factors.

These factors are usually inter-dependent so that rather than a chain of causes, there is usually a causal web. These inter-relationships should be considered when selecting a species for aquaculture, choosing a site and designing grow-out facilities and management practices.

### **Avoid stress**

Appropriate growing conditions that do not stress the fish are essential for successful culture. While species can tolerate sub-optimal conditions for short periods (e.g. for prawns, short periods of low temperature, dissolved oxygen levels or salinity levels), they will become stressed with reduced FCRs and greater susceptibility to disease than species better suited to those conditions.

Many of the potential disease parasites, viruses and bacteria occur naturally in the environments and may be present in the ponds or tanks. The natural defence systems of healthy fish will ward off infection. However, once a fish becomes stressed, opportunity for the disease to cause infection increases via the lowering of the immune systems. The stress may also occur during handling (e.g. grading, harvesting, transferring between ponds, transport to market), heavy predation from predators (such as cormorants), drug treatments, poor water quality, undernourishment or overcrowding.

Prevention of stress is fundamental to maintaining disease free stock. Good husbandry techniques for disease prevention include:

- maintenance of appropriate water quality standards;
- maintenance of optimum stocking rates to prevent overcrowding;
- quarantine management for new stock before introduction into the ponds;
- development of feeding regimes to maintain water quality;
- regular inspection of fish should be undertaken to monitor health and implement of disease management protocols if necessary.

### **Disease management protocols**

Some diseases are not naturally occurring and every effort should be taken to ensure that these are not introduced onto the farm from hatcheries, other farms or the wider environment. Measures should include stock protocols, worker protocols and in some circumstances exclusion of birds. A useful website on this subject is [www.dah.csiro.gov.au](http://www.dah.csiro.gov.au). Many of the disease outbreaks (if caught early) are easily treatable, with little or no effect on the productivity of the crop. However, some fish diseases are difficult to treat and cause widespread mortality in the aquaculture facilities as well as in the wild. Disease in hatcheries can be a particular problem, both in terms of losses in the hatcheries and also through putting at risk customer farms or native fish stocks (through restocking programs).

Good practice for disease management should include:

- At the first signs of disease outbreak specimens should be removed and analysed for disease symptoms and water quality should be checked. If no obvious manageable causes are found then a qualified veterinarian should be consulted immediately.
- Appropriate training for all staff in health management and disease recognition but in particular key personnel who should have specialised training in the recognition and diagnosis of common fish diseases. These personnel should be made aware of the importance of keeping up-to-date with the latest disease identification and management.
- Appropriate equipment on site to assist in the accurate diagnosis of diseases including a microscope and testing kits.
- Appropriate protocols to prevent the inadvertent introduction of disease onto the farm from stock movement, employees, visitors or equipment.
- Appropriate protocols to deal with disease notification and management should an outbreak occur.

## **DISEASE PROFILES**

### ***Abalone***

Abalone appears to have one major disease known as Perkinsus which naturally occurs predominantly from Jervis Bay and north along the coastline. It can cause devastating losses to stock and is linked to high summer water temperatures and is not treatable.

Abalone are also susceptible to Mud worm and Vibrio bacterial infections.

### ***Prawns***

To protect native prawn species and reduce the risk of disease introduction, no live penaeids are allowed into Australia. So far, this appears to have prevented the introduction of major penaeid viral diseases such as infectious hypodermal-haematopoietic necrosis virus disease (IHHNV), yellow-head and white spot virus, which has caused very serious problems in countries that have allowed importations. Other viral diseases have occurred in Australia (for example Baculovirus), although this only appears to be a problem in hatcheries cultured prawns under sub-optimum conditions. Although bacterial, fungal and protozoan infections have been recorded in Australian prawn hatcheries and ponds, very few have caused serious problems.

### ***Freshwater Crayfish***

Australian freshwater crayfish appear to have only one major disease, the so-called porcelain or white tail disease, caused by a microsporidian. The disease appears to be transmitted through cannibalism of dead individuals. The disease cannot be treated, but can be managed in aquaculture if stocks are periodically examined, and diseased animals removed. Microsporidians have been found in all three *Cherax* species.

### ***Silver and Golden Perch***

Infection with disease and parasites is a response by perch to stressors occurring in their environment. The most common of these would be deterioration of water quality. Most pathogens and parasites occur naturally in ponds and natural aquatic environments, however the natural defence systems of healthy fish will help ward off infection. Once fish become stressed with a lowering of its immunity defences, opportunity for the disease to infect increases. Infection may also occur when fish are handled e.g. grading, harvesting, moving between ponds, or stress caused by heavy presence of predators such as cormorants, or from inappropriately high stocking densities.

Prevention is the best cure, and a combination of good husbandry and management techniques will ensure that stock remains relatively disease free. General husbandry should include quarantine of all new stock to the site, with regular salt baths of new batches at a concentration of 5-10ppt for at least 1 hour before placement into any of the nursery ponds. Optimum water quality should be maintained to relieve stress on the fish, and regular inspections to monitor health. At the first signs of a disease outbreak a sample of fish should be removed and analysed for obvious disease

symptoms, and water quality should be checked. If a source of infection or disease cannot be identified, a qualified fish veterinarian should be consulted. Most fungal and parasitic infections can be treated with a combination of salt baths and formalin.

### **Trout**

Trout are vulnerable to disease if stressed. Temperature stress is the most common problem in NSW, then overcrowding and low oxygen. Common parasites include Ichthyophthirius ('Ich' or 'whitespot') and Trichodina, both protozoans. Poor hygiene is also a common cause of bacterial disease in hatcheries, where any disease can spread rapidly if not identified and treated. Introduced trout diseases in the river stocks are a source of concern. In particular, outbreaks of trout EHN virus have resulted in new quarantine regulations.

### **Murray Cod**

Like any fish, Murray Cod can be susceptible to various pathogens and diseases. Disease outbreaks are often in response to stressors being placed on the stock, which in most cases could be avoided. Prevention is the best cure, and a combination of good husbandry and management techniques will ensure that stock remains relatively disease free. General husbandry should include quarantine of all new stocks to the site, with regular salt baths of new batches at a concentration of 5-10ppt for at least 1-hour before placement into any of the grow-out tanks. Water quality should be maintained to relieve stress on the fish, and regular inspection of fish should be undertaken to monitor health. Upon the first signs of a disease outbreak a sample of fish should be inspected for obvious disease symptoms, and water quality should be checked. If a source of infection or disease cannot be identified, a qualified fish veterinarian should be consulted.

### **Barramundi**

Barramundi are naturally susceptible to most bacterial, fungal and parasitic infections, particularly at times of stress. This can usually be avoided by appropriately quarantining new stock before release into culture tanks, maintaining water quality and a stress free environment, and regular disease monitoring of stock. In the event of disease outbreak, stock can sometimes be effectively treated by salt or freshwater baths, or via veterinarian prescribed medicines. The barramundi is known to be a potential carrier of the barramundi Encephalitis Virus (BEV) which has potential to affect a number of species native to NSW. As such Department of Primary Industries has imposed tight restrictions on the culture of barramundi in this State. To address these concerns Department of Primary Industries have included provision within the Barramundi Farming Policy for the sterilisation of all effluent to be removed, as well as a specific protocol for the importation of barramundi fingerlings from out of state. This includes the testing of fingerlings for presence of the BEV as well as other diseases and virus. This policy will not only reduce the chance of translocating the virus to NSW, but also ensures that the farmer has a guarantee of healthy good quality seedstock.

### **Eels**

Because glass eels are sourced from the wild, they may already be infested with various parasites or diseases, or alternatively act as disease carriers. However, the acclimatisation to freshwater that usually occurs after capture eliminates most parasitic infections through osmotic forces. Bacterial and fungal infections on eels raised in freshwater can usually be treated by simple salt baths at a rate of around 10ppt. Eels, like most fish, are particularly susceptible to infection at times of high stress, for example when water quality has deteriorated, there is constant presence of predators in ponds, or at times of harvesting and handling. While eels are generally quite tough, they still need to be handled with care. As in most forms of aquaculture, the best cure is prevention and a combination of good husbandry and management techniques will ensure that stock remains relatively disease free. The maintenance of disease free stock involves the quarantining of all new eels to the site, and regular salt bathing of eels in quarantine facilities. Water quality should be maintained to relieve stress, and regular inspection of eels or elvers should be undertaken to monitor health and condition. At the first signs of a disease outbreak a sample of eels should be inspected for obvious disease symptoms, and water quality should be monitored. If a source of infection or disease cannot be identified, a professional fish veterinarian should be consulted.

## Reference Sources

### ***Barramundi***

- Barlow, C 1998 'Barramundi'. The New Rural Industries - A Handbook for Farmers and Investors. Rural Industries Research and Development Corporation.
- NSW Fisheries 1997 Barramundi Farming Policy - NSW Fisheries Policy Paper.
- Phillips, C 1998 'Barramundi Farming'. Proceedings from the Queensland Warmwater Aquaculture Conference (Status and Potential) 1998. Aquaculture Information Technologies (ed).
- Rimmer, M 1995 Barramundi Farming, An Introduction. Queensland Department of Primary Industries.

### ***Eels***

- Beumer, J 1983 'Suitability of *Anguilla australis* for Intensive Culture'. Aust Aqua, (4)2:13-18.
- Forteach, N 1994 'Eel Culture In Australia'. Austasia Aquaculture, 8(5):54-55.
- WBM Oceanics Australia 1995 'Feasibility Study for Eel Aquaculture in Queensland. Queensland Department of the Premier.
- Wray, T 1995 'Eels in Japan'. Fish Farming International. 22(10):10-11.
- NSW Fisheries 1998 'Eel Aquaculture Policy'.
- Reeve, J 1998 'Eel Farming'. Proceedings from the Queensland Warmwater Aquaculture Conference (Status and Potential) 1998. Aquaculture Information Technologies (ed).

### ***Freshwater Crayfish***

- Jones, C.M. 1990 The Biology and Aquaculture Potential of the Tropical Freshwater Crayfish. *Cherax quadricarinatus*. Queensland Department of Primary Industries Information Series Q190028.
- Mills, B.J. 1989 Australian Freshwater Crayfish. Handbook of Aquaculture. Freshwater – Crayfish Aquaculture, Research and Management, RSD. 778 Lymington, Tasmania 7109.
- Morrissy, N.M. 1981 Marron and Marron Farming. Department of Fisheries and Wildlife. Western Australia.
- NSW Fisheries 1994 Freshwater Crayfish Advisory Pack. NSW Fisheries, Sydney.
- Shelly, C.C. & Pearce, M.C. 1990 Farming the Redclaw Freshwater Crayfish. Northern Territory Department of Primary Industry and Fisheries Fishery Report No. 21.

### ***Murray Cod***

- Mosig, J 1999 'Murray Cod Grower Finds Success in Tanks'. Austasia Aquaculture, 13(1):14-16.

### ***Trout***

- Roberts, R. J. and Shepherd, C. J. 1974 Handbook of Trout and Salmon Diseases. Fishing News Books.
- Rowland S. J. and Ingram, B. A. 1991 Diseases of Australian native freshwater fishes. Fisheries Bulletin No. 4. NSW Agriculture & Fisheries.
- Stevenson, J. P. 1980 Trout Farming Manual. Fishing News Books

### **Silver Perch**

- Cacho, O., H. Kinnucan and S. Sindelar 1986 Catfish farming risks in Alabama. Alabama Agricultural Experiment Station Circular 327, Auburn, Alabama, USA
- Lambert, N 1998 'Commercial Aquaculture of Silver Perch'. Proceedings from the Queensland Warmwater Aquaculture Conference (Status and Potential) 1998. Aquaculture Information Technologies (ed).
- Mifsud, C 1998 'Silver Perch Makes a Splash'. Fisheries of NSW – The Journal of Sustainable Fishing. NSW Fisheries.
- NOFARIC 1996 Marketing Silver Perch (*Bidyanus bidyanus*) Prepared by Ruello and Associates Fisheries Consultants. NOFARIC, Sydney, Australia.
- NSW Fisheries 1994 Silver Perch Fish Farming Advisory Package.
- NSW Fisheries 1994 Aquaculture Permits for Silver Perch – NSW Fisheries Policy Paper.
- Rowland, S and Bryant, C (eds) 1994 Silver Perch Culture – Proceedings of Silver Perch Aquaculture Workshops, Grafton and Narrandera, April 1994. Austasia Aquaculture Publication.
- Rowland, S 1998 'Silver Perch'. The New Rural Industries – A Handbook for Farmers and Investors. Rural Industries Research and Development Corporation.
- Van der Ploeg, M., and C. E. Boyd, 1991 Geosmin production by cyanobacteria (blue-green algae) in fish ponds at Auburn, Alabama. Journal of the World Aquaculture Society 22:4 (204-216).

### **Prawn**

- Akiyama, D. M. (Editor) 1990 Proceedings of the Southeast Asia Shrimp Farm Management Workshop. American Soybean Association, Singapore.
- Boyd, C. E. 1989 Water Quality Management and Aeration in Shrimp Farming. Fisheries and Allied Aquacultures Departmental Series No. 2, Alabama Agricultural Experimental Station, Auburn
- Fast, W.W. & Lester, L.J. (Ed) 1992 Marine Shrimp Culture: Principles and Practices. Elsevier, N.Y.
- Fast, A. W & Lester, L. J. (Ed) 1992 Marine Shrimp Culture: Principles and Practices. Elsevier, N.Y
- Macleay, J. I., Dizon, L. B. & Hosillos, L. V. (Ed) 1986 Proceedings of the First Fisheries Forum. The Asian Fisheries Society, Manila, Philippines.
- McVey, J. P. 1983 CRC Handbook of Mariculture. Volume 1, Crustacean Aquaculture. CRC Press Inc. Boca Raton, Florida.
- NOFARIC 1995 Biology and farming of the kuruma prawn (*Penaeus japonicus*). Prepared by the Pacific Seafood Management Consulting Group Pty. Ltd. NOFARIC, Sydney, Australia.