

Assessment Authors and Year

Smoothey, AF 2021. NSW Stock Status Summary 2021/22 – Blue-eye Trevalla (*Hyperoglyphe antarctica*). NSW, Department of Primary Industries, Fisheries. 16 pp.

Stock Status

Current stock status	On the basis of the evidence contained within this assessment, Blue-eye Trevalla are currently assessed as sustainable .
----------------------	---

Stock structure & distribution

Blue-eye Trevalla (*Hyperoglyphe antarctica*) are benthic species that are found in areas of rocky ground on continental slopes waters to depths of 200-900 m. They are distributed in continental slope waters off South America, South Africa, New Zealand and Australia. Their Australian distribution stretches along the southern continental margin in waters from Moreton Island in Queensland to 30°S in Western Australia. Blue-eye Trevalla also occur on the seamounts off eastern Australia and south of Tasmania, Lord Howe Island and Norfolk Island. Adults and sub-adults occur in mid-water at depths of around 500 m and are associated with rocky ground on the continental slope where the majority of fish are found between 200 and 600 m, but a small number have been reported to occur at depths of up to 900 m. They generally remain close to the seabed during the day and move into the water column at night.

In recent years, stock structuring has been reported based on phenotypic variation in age and growth, otolith chemistry and potential larval dispersal between regions suggests spatial patterns may delineate natural subpopulations of Blue-eye Trevalla (Williams et al. 2017). Four geographically distinct subpopulations around the Australian coast were identified: 'West' – comprising continental slope fishing grounds off Western Australia, South Australia and western Victoria to western Tasmania; 'South' – continental slope grounds around Tasmania and north eastwards to eastern Bass Strait; 'East' – fishing grounds on the NSW continental slope and Tasmanian seamounts; and 'Offshore' – fishing grounds on the Lord Howe Rise (Williams et al. 2017). Blue-eye Trevalla stock areas do not reflect truly separated biological stocks because there is some exchange between them during pelagic early life history (Williams et al. 2017). However, local-scale residency by adults implies there are discrete adult populations on the continental slope and seamounts and that there is not extensive migration between them. Within the Commonwealth, Blue-eye Trevalla are managed as a single biological stock in the Southern and Eastern Scalefish and Shark Fishery (SESSF; Patterson et al. 2021), however, these findings led to separate RBCs being determined for the slope and seamount stocks, but a global TAC applied (AFMA 2018a) and catch restrictions introduced for the seamount stock for the 2019/20 fishing season.

Scope of this assessment

The fishery scientific assessment summarised in this report and considered adequate to meet the legislative requirements for supporting a Total Allowable Catch (TAC) determination for NSW Blue-eye Trevalla is that done by the CSIRO, commissioned by the Australian Fisheries Management Authority (AFMA) and published as 'Blue-eye Trevalla (*Hyperoglyphe antarctica*)' by the Australian

Bureau of Agricultural and Resource Economics and Sciences (Patterson et al. 2021; hereinafter referred to as the Commonwealth assessment).

Assessment of the status of the stock of Blue-eye Trevalla that is fished by commercial and recreational fishers in New South Wales (NSW) is principally based on the modelling and assessment done for this species by the Commonwealth of Australia for Eastern Australia. The Commonwealth fisheries, primarily Southern and Eastern Scalefish and Shark Fishery (SESSF) takes 85-95% of the historical catch of Blue-eye Trevalla. The primary mechanism for controlling the harvest of Blue-eye Trevalla in SESSF is through the allocation of an Annual Total Allowable Catch (TAC). Determination of annual TACs for the Commonwealth SESSF is based on the SESSF Harvest Strategy Framework (HSF) (AFMA, 2017) that derives from the Commonwealth Fisheries Harvest Strategy Policy (HSP) (DAFF, 2007).

The Commonwealth assessment for Blue-eye Trevalla is done as a Tier 4 assessment on catch and CPUE for the slope stock and as a Tier 5 for the seamount stock. The Tier 4 analysis determines an Recommended Biological Catch (RBC) by selecting CPUE reference points that are taken as proxies for the estimated BLIM and BTARG. This is done by assuming that the CPUE is proportional to stock abundance. If the stock was at unexploited equilibrium at the start of fishing, then the initial CPUE level at the start of the time series would correspond to the unexploited biomass or B₀, and the other reference points are the appropriate fractions of this (e.g. 20% for B₂₀). The Tier 5 assessment for Blue-eye Trevalla uses catch at maximum sustainable yield (MSY) and age-structured stock reduction analysis approaches to inform the TAC setting.

This assessment of the status of Blue-eye Trevalla, in waters under NSW jurisdiction, comprises:

- (1) a summary of the most recent Commonwealth stock assessment for Blue-eye Trevalla and current determinations of status based on criteria specified by the Commonwealth and also those used for the Status of Australian Fish Stocks;
- (2) the rationale by which the Commonwealth assessment for Blue-eye Trevalla is considered to be relevant and valid for determining the status of the Blue-eye Trevalla stock fished within NSW jurisdiction (Appendix 1);
- (3) information that may inform the determination of the 2022-23 NSW TAC for Blue-eye Trevalla in the Ocean Trap and Line – Line East Fishery (Appendix 2). This is done in the absence of: (i) a formal NSW harvest strategy for this species/fishery; and (ii) a formal resource sharing agreement between NSW and the Commonwealth.

Biology

Blue-eye Trevalla are commonly found around 60 cm in length and 3 kg, however, they can grow up to 1.4 m in length and 50 kg, with maximum age reported to be 76 years. Female Blue-eye Trevalla reach sexual maturity at 11-12 years of age (72 cm fork length), while males mature at 8-9 years of age (62 cm fork length). Spawning occurs in summer and autumn in waters from central New South Wales to north-eastern Tasmania. Eggs are released in batches of 3-4 and females produce 2-11 million eggs per spawning season (AFMA, 2021a).

FISHERY STATISTICS

Catch information

Commercial


Fishery statistics presented in this report are restricted to those used to inform the Commonwealth assessment and are summarised here from Patterson et al. (2021) and references therein. The Commonwealth assessment of Blue-eye Trevalla uses data from the Commonwealth Ocean Trawl Fishery within Commonwealth fishing zones 10 and 83 (south-eastern Australia).

Descriptions of NSW fishery statistics are provided in Appendix 2 of this report and the changes in NSW commercial fishery reporting requirements and sources of NSW commercial fishery data are discussed.

The description of catch information below is summarised from Paterson et al. (2021).

In 1997 catches of Blue-eye Trevalla peaked at over 800 t and gradually declined since then (Figure 1). Commonwealth landed catch in the 2016/17 fishing season was 432 t, 215.5 t in the 2019/20 fishing season and in 2020/21 was 224.6 t. The weighted average discards and state catches over the past four calendar years (2016 to 2019) were 7 t and 22.1 t, respectively (Althaus et al. 2020). Commonwealth catches have varied in response to changes in the TAC, but in some years, there has been uncaught quota.

Figure 1. Blue-eye Trevalla annual catches (CTS, SHS and states) and fishing season TACs, 2006/07 to 2020/21 (Source: AFMA catch disposal records, cited in Patterson et al. 2021).



Recreational & Charter boat

Recreational catches have not been accounted for in the Commonwealth assessment of Blue-eye Trevalla. The model is conditioned on commercial catch data alone. Accounting for recreational catch has been raised as an issue for consideration in Commonwealth assessments (SESSF RAG 2017). In NSW, there is a combined recreational bag limit of five Hapuku, Banded Rockcod, Bass Groper, Gemfish and Blue-eye Trevalla. Recreational catch of Blue-eye Trevalla in New South Wales are unknown. Surveys of the catches in NSW by NSW-resident recreational fishers during

2013-14 and by 1-3 year licence holders in 2017/18 did not report any catches of Blue-eye Trevalla (West et al, 2015; Murphy et al., 2020). There is, however, anecdotal evidence, including photographs and reports on social media websites, indicating that Blue-eye Trevalla are targeted and captured by a small subset of recreational anglers using specialist gear. State-wide operators within the nearshore charter fishery landed 110 Blue-eye Trevalla during the 2017/18 survey period, with 58% caught during summer (Hughes et al. 2021). The interpretation of the survey results and the evidence from social media is that catches of Blue-eye Trevalla by recreational fishers in NSW are negligible, relative to the magnitude of commercial catches.

Indigenous

Neither the Commonwealth assessment or the process for determining the Commonwealth TAC from RBC includes estimates of Blue-eye Trevalla catches by the Aboriginal sector. Given the depths inhabited by Blue-eye Trevalla, this species would not have been a traditional target species for Aboriginal fishers. As for the recreational catch, any catches by the Aboriginal fishers in recent history are negligible, relative to the magnitude of commercial catches. Thus, the omission of Aboriginal catch from the model and Commonwealth assessment has little effect on the assessment outcome of the Blue-eye Trevalla stock.

Illegal, Unregulated and Unreported

The level of Illegal Unregulated and Unreported (IUU) fishing has not been quantified.


Fishing effort information

N/A

Catch Rate information

Standardised CPUE (0.53) for the slope stock has declined over the last five years from above the target reference point in 2014 to a point just above the limit reference point (0.51) in 2019 (Figure 2). Two factors that could influence the catch rates and fishing behaviour resulting in a low bias for CPUE are depredation by killer whales (orcas, *Orcinus orca*) and exclusions from historical fishing grounds following closures implemented to rebuild the stocks of gulper sharks (Patterson et al. 2021).

Figure 2. Standardised auto-longline and dropline CPUE index for Blue-eye Trevalla to the east and west of Tasmania, 1997 to 2019 (Source: Sporcic 2018, cited in Patterson et al. 2021).



STOCK ASSESSMENT

The management of the stock for the 2020/21 fishing season was based on analyses done by Sporcic (2018, 2020) and Haddon and Sporcic (2018a, b) and summarised from Patterson et al. (2021). Based on the recent evidence of stock structuring (Williams et al. 2017), the 2018 analysis split the stock into two regions (slope and seamount populations), with each analysed separately to inform the determination of a recommended biological catch for the 2020/21 fishing season. The Commonwealth SESSF HSF assessment (AFMA 2018a, b, c) for Blue-eye Trevalla is a Tier 4 analysis for the slope stock and Tier 5 analysis for the seamount stock (due to unreliable catch-per-unit-effort [CPUE] data) (AFMA 2020a). Although the 2018 Tier 4 analysis (Sporcic 2018) informed the management of the slope stock for the 2020/21 fishing season, a new Tier 4 analysis was completed in 2020 (Sporcic 2020).

Stock Assessment Methodology

Year of most recent assessment:

2020 - slope stock (Sporcic 2018, 2020).

2018 - seamount stock (Haddon and Sporcic 2018a, b; Sporcic 2018).

Assessment method:

Commonwealth Tier 4 analysis - slope stock - standardised CPUE (including discards).

Commonwealth Tier 5 analysis - seamount stock - catch at maximum sustainable yield (MSY) and age-structured stock reduction analysis approaches.

Main data inputs:

CPUE - Commonwealth Trawl Fishery; Zones 20-50; Depth 200-600 m.

Stock Status Summary – 2021/22



NSW Stock Status Summary – Blue-eye Trevalla (*Hyperoglyphe antarctica*)

Catch (Total) is the sum of Discards, State (Vic, Tas and NSW), Non-trawl and SEF2 catches (Haddon and Sporcic 2018a, b).

Discard rates.

Commonwealth Tier 5 - catch at maximum sustainable yield (MSY) and age-structured stock reduction analysis approaches.

Key model structure & assumptions:

Tier 4 – Standardised CPUE (Commonwealth harvest strategy policy).

Assumptions (see Haddon 2016): catch rate provides a relative index of abundance (not subject to hyper-stability or hyper-depletion and not overly influenced by other factors not accounted for through standardisation); the reference period provides a good estimate of the stock when at a depletion level of $0.48B_0$; estimates of catch during the target period are accurate.

Sources of uncertainty evaluated:

Uncertainty associated with Tier 4 assessment (see Haddon 2016): factors considered in the CPUE standardisation: Year, Vessel, Month, Zone, Depth category and Month:Zone; investigation of additional zones (84 and 85). Two factors that could influence catch rates and fishing behaviour, resulting in a low bias for CPUE, include the presence of killer whales (orcas—*Orcinus orca*) and Commonwealth fishery closures implemented to rebuild stocks of gulper sharks. The previous analysis by Haddon (2016) did not detect large effects on CPUE due to the closures, but uncertainty remains about the effect of killer whale depredation on CPUE (Patterson et al. 2021).

Status Indicators - Limit & Target Reference Levels

Biomass indicator or proxy	Standardised CPUE (AFMA 2017)
Biomass Limit Reference Point	Standardised CPUE at $0.20B_0$ = Limit Reference Point
Biomass Target Reference Point	Standardised CPUE at $0.48B_0$ = Target Reference Point
Fishing mortality indicator or proxy	Catch (including discards) as a proportion of RBC. Note, the RBC calculation does not account for predicted discards of predicted State catches.
Fishing mortality Limit Reference Point	NA
Fishing Mortality Target Reference Point	CPUE ₄₈ at or above target, F_{48} (Fishing mortality rate that achieved B_{48})

Stock Assessment Results

The Tier 4 slope stock analysis (Sporcic 2018) suggested that the previous steep decline in CPUE (2013 to 2016) had levelled out and remained between the target and limit reference points as defined by the SESSF HSF (AFMA 2017). These analyses produced an RBC of 439 t. The South

Stock Status Summary – 2021/22



NSW Stock Status Summary – Blue-eye Trevalla (*Hyperoglyphe antarctica*)

East Resource Assessment Group (SERAG) recommend using the RBC to set the second year of a 3-year MYTAC for the 2020/21 fishing season. The 2020 Tier 4 slope analysis (Sporcic 2020) revealed, however, a decline in the most recent (2019) CPUE to just above the limit reference point (Figure 1). These analyses produced an RBC 227t for the 2021/22 fishing season. Industry observed poor catches in the 2019/20 fishing season (AFMA 2020e) and SERAG noted that catches needed to be reduced in accordance with the RBC to limit the decline (AFMA 2021b).

The Tier 5 age-structured stock reduction analysis of the seamount population predicted that constant catches of around 25 t for lower-productivity scenarios and 48 t for higher-productivity scenarios would lead to relative stability in depletion (Haddon & Sporcic 2018b). A maximum sustainable yield (MSY) analysis of the seamount catch resulted in an MSY of around 46-50 t, with a depletion estimate of about 33% of the unfished biomass (Haddon & Sporcic 2018), however, this needs to be interpreted with caution due to the high uncertainty. It was predicted, based on the catch MSY, that constant catches of 40 t or less would lead to relative stability in depletion (AFMA 2018c, d). The application of the SESSF Tier 4 harvest control rule to the outputs of the standardised CPUE series for the slope stock generated a single-year RBC of 439 t. The South East Resource Assessment Group (SERAG) agreed to an RBC of 36 t for the seamount stock, based on the output of the age-structured stock reduction analysis and catch-MSY analysis for the 2020/21 fishing season (AFMA 2018c, d). When combined with the RBC of 439 t for the slope stock, this led to a total combined RBC of 475 t for the 2020/21 fishing season.

SERAG have recommended that the Tier 4 seamount and Tier 5 slope stock analyses be updated in 2021 (AFMA 2020d) due to decline in the recent (2019) CPUE from the slope stock and the uncertainty in the 2018 analysis for the seamount stock.

For the 2020/21 fishing season, the agreed TAC was 448 t, and the recommended biological catch (RBC) for the slope stock was 439 t, while the RBC for the seamount was 36 t. The catch and discards combined were estimated to be 253.7 t (97.3% from SSH; 2.7% CHS), which is below the combined RBC of 475 t (Figure 1). This suggests that the fishing mortality in 2020/21 would be unlikely to deplete the stock to a level below its biomass limit reference point. The stock is therefore classified as **not subject to overfishing**.

The 2020 slope stock analysis (Sporcic 2020) predicted that the recent average standardised CPUE was between the target reference point and limit reference point, and the 2018 seamount analysis (Haddon & Sporcic 2018a, b) estimated that constant catches of 40 t or less would see biomass maintained at around 0.33B₀ for the seamount stock (Figure 2). The Blue-eye Trevalla stock is, therefore, classified as **not overfished**.

The status of the eastern Australian stock of Blue-eye Trevalla was defined as **sustainable**, under the criteria for SAFS in 2016 (the first year of SAFS assessment; Georgeson et al. 2016), 2018 and 2020 (Emery et al. 2020).

Stock Assessment Result Summary

Biomass status in relation to Limit	Recent average standardised CPUE (Biomass proxy) is between the Target and Limit Reference Point (Sporcic 2018, Patterson et al. 2021)
Biomass status in relation to Target	Biomass proxy (Standardised CPUE) is above Target Reference Point (Patterson et al. 2021).

Stock Status Summary – 2021/22



NSW Stock Status Summary – Blue-eye Trevalla (*Hyperoglyphe antarctica*)

Fishing mortality in relation to Limit	N/A
Fishing mortality in relation to Target	N/A
Current SAFS stock status	Sustainable
Current Commonwealth stock status	Sustainable

Fishery interactions

Interactions between the Commonwealth Trawl and Auto-longlining Fisheries are described by Haddon and Sporcic (2018a, b), who associate declines in the trawl sector since the mid-2000s with increased catches in the auto-longlining sector.

Commonwealth fisheries interact with other commercial and non-commercial bycatch and discard marine species, a range of endangered threatened and/or protected species and marine habitats (AFMA 2014, Wayte et al. 2007).

Qualifying Comments

Supplementary information relevant for to the interpretation of the assessment is provided in Appendix 1 and 2.

References

- AFMA, 2017. Harvest strategy framework for the Southern and Eastern Scalefish and Shark Fishery 2009 (amended March 2017). Australian Fisheries Management Authority, Canberra. www.afma.gov.au/wp-content/uploads/2017/03/SESSF-Harvest-Strategy-Framework-2017-final.pdf
- AFMA, 2018a. Total allowable catch recommendations for Southern and Eastern Scalefish and Shark Fishery (SESSF) 2019-20 fishing year, Australian Fisheries Management Authority, Canberra.
- AFMA, 2018b. Southern and Eastern Scalefish and Shark Fishery (SESSF) South East Resource Assessment Group (SERAG), minutes, 19-21 September 2018, Hobart, Australian Fisheries Management Authority, Canberra.
- AFMA, 2018c. Species summaries for the Southern and Eastern Scalefish and Shark Fishery (SESSF): for stock assessments completed in 2018 in preparation for the 2019-20 fishing seasons, Australian Fisheries Management Authority, Canberra.
- AFMA, 2020a. Harvest strategy framework for the Southern and Eastern Scalefish and Shark Fishery (SESSF) 2009: amended (2020), Australian Fisheries Management Authority, Canberra.
- AFMA, 2020b. 'Southern and Eastern Scalefish and Shark Fishery (SESSF) Resource Assessment Group (SESSFRAG) data meeting, minutes, 25–26 August 2020', Australian Fisheries Management Authority, Canberra.
- AFMA, 2020c. Total allowable catch recommendations for the Southern and Eastern Scalefish and Shark Fishery (SESSF) 2020–21 fishing year: advice to SEMAC, Australian Fisheries Management Authority, Canberra.

Stock Status Summary – 2021/22



NSW Stock Status Summary – Blue-eye Trevalla (*Hyperoglyphe antarctica*)

- AFMA, 2020d. 'Southern and Eastern Scalefish and Shark Fishery (SESSF) South East Resource Assessment Group (SERAG) meeting 2.2, minutes, 9–10 December, 2020', Australian Fisheries Management Authority, Canberra.
- AFMA, 2020e. 'Southern and Eastern Scalefish and Shark Fishery South East Management Advisory Committee (SEMAC) meeting, minutes, 4–6 February 2020', Australian Fisheries Management Authority, Canberra.
- AFMA, 2021a. Blue-eye Trevalla. Australian Fisheries Management Agency.
<https://www.afma.gov.au/fisheries-management/species/blue-eye-trevalla>
- AFMA, 2021b. *Southern and Eastern Scalefish and Shark Fishery (SESSF) species summaries 2021*, Australian Fisheries Management Authority, Canberra.
- Althaus, F, Burch, P and Thomson, R 2020. Southern and Eastern Scalefish and Shark Fishery (SESSF) catches and discards for TAC purposes using data until 2019, CSIRO Oceans and Atmosphere, Hobart.
- Commonwealth of Australia, 2007. Commonwealth fisheries harvest strategy policy, Department of Agriculture and Water Resources, Canberra.
www.agriculture.gov.au/SiteCollectionDocuments/fisheries/domestic/hsp.pdf
- Commonwealth of Australia, 2017. Commonwealth fisheries harvest strategy policy, Department of Agriculture and Water Resources, Canberra.
www.agriculture.gov.au/SiteCollectionDocuments/fisheries/domestic/consultation-draft-harvest-strategy-policy.pdf
- DAAF, 2007. Commonwealth Fisheries Harvest Strategy: policy and guidelines, Australian Government Department of Agriculture, Fisheries and Forestry, Canberra.
- Emery, T., Roelofs, A., Smoothey, AF., Trinnie, F., Wakefield, C. and Lyle, J, 2020. Blue-eye Trevalla *Hyperoglyphe antarctica*, In: Toby Piddocke, Crispian Ashby, Klaas Hartmann, Alex Hesp, Patrick Hone, Joanne Klemke, Stephen Mayfield, Anthony Roelofs, Thor Saunders, John Stewart, Brent Wise and James Woodhams (eds) 2021, *Status of Australian fish stocks reports 2020*, Fisheries Research and Development Corporation, Canberra.
- Georgeson, L., Roelofs, A., Wakefield, C., Lyle, J. and R. Chick, 2016. Blue-eye Trevalla *Hyperoglyphe antarctica*, In: Stewardson, C., Andrews, J., Ashby, C., Haddon, M., Hartmann, K., Hone, P., Horvat, P., Mayfield, S., Roelofs, A., Sainsbury, K., Saunders, T., Stewart, J., Stobutzki I. and B. Wise, Eds. 2016. *Status of Australian Fish Stocks reports 2016*. Fisheries Research and Development Corporation, Canberra.
www.fish.gov.au/report/16-Blue-eye-Trevalla-2016
- Haddon, M. 2016. Blue-Eye (*Hyperoglyphe antarctica*) Tier 4 Analysis using Catch-per-Hook for Auto-Line and Drop-Line from 1997–2015. CSIRO Oceans and Atmosphere, Hobart. 25p.
- Haddon, M and M. Sporcic, 2017a. Statistical CPUE standardisations for selected SESSF species (data to 2016), CSIRO Oceans and Atmosphere, Hobart.
- Haddon, M and M. Sporcic, 2017b. Tier 4 assessments for selected SESSF species (data to 2016), CSIRO Oceans and Atmosphere, Hobart.
- Haddon, M and M. Sporcic, 2018a. Draft blue-eye trevalla tier 5 eastern seamount assessment: catch-MSY analysis, CSIRO Oceans and Atmosphere, Hobart.
- Haddon, M and M. Sporcic 2018b. Draft tier 5 blue-eye trevalla eastern seamount assessment: age-structured stock reduction analysis, CSIRO Oceans and Atmosphere, Hobart.

- Henry, G.W. and J. M. Lyle. 2003. The national recreational and Indigenous fishing survey, Fisheries Research and Development Corporation, Canberra.
- Hughes, J.M, Johnson, D, Murphy, J.J and Ochwada-Doyle, F.A, 2021. The NSW Recreational Fisheries Monitoring Program – Charter Fishery Monitoring 2017/18. NSW DPI – Fisheries Final Report Series No. 159. ISSN 2204-8669. pp 97.
- Macbeth, W. G. and C. A. Gray. 2015. Observer-based study of commercial line fishing in waters off New South Wales, NSW DPI – Fisheries Final Report Series No. 148. Commercial Fishing Trust Fund Project no. FSC2006/179. NSW DPI, Coffs Harbour.
- Murphy, J.J., Ochwada-Doyle, F.A., West, L.D., Stark, K.E. and Hughes, J.M., 2020. The NSW Recreational Fisheries Monitoring Program - survey of recreational fishing, 2017/18. NSW DPI - Fisheries Final Report Series No. 158.
- Patterson, H, Bromhead, D, Galeano, D, Larcombe, J, Woodhams, J and R, Curtotti, 2021. Fishery status reports 2021, Australian Bureau of Agricultural and Resource Economics and Sciences, Canberra. CC BY 4.0. <https://doi.org/10.25814/5vahf-ng93>
- SESSF RAG (Southern and Eastern Scalefish and Shark Fishery Resource Assessment Group), 2017. Chair's Meeting Minutes 16–17 March 2017. www.afma.gov.au/wp-content/uploads/2017/05/2017-SESSFRAG-Chairs-Meeting-Minutes-FINAL.pdf
- Sporcic, M. and M. Haddon, 2016. Catch rate standardizations for selected SESSF species (data to 2015). CSIRO Oceans and Atmosphere Flagship, Hobart. 214p.
- Sporcic, M, 2018. Tier 4 assessments for selected SESSF species (data to 2017), CSIRO Oceans and Atmosphere, Hobart. 2018
- Sporcic, M, 2020. Draft tier 4 assessments for selected SESSF species (data to 2019), CSIRO Oceans and Atmosphere, Hobart.
- Suthers, I. M., Young, J. W., Baird, M. E., Roughan, M., Everett, J. D., Brassington, G. B., Byrne M., Condie, S. A., Hartog, J. R., Hassler, C. S., Hobday, A. J., Holbrook, N. J., Malcolm H. A., Oke, P. R., Thompson, P. A., and K. Ridgway, 2011. The strengthening East Australian Current, its eddies and biological effects—an introduction and overview. Deep Sea Research Part II: Topical Studies in Oceanography 58:538–546.
- Thomson, R and J. Upston, 2016. SESSF catches and discards for TAC purposes, 9 November 2016, report prepared by CSIRO Wealth from Oceans Flagship for AFMA, Canberra.
- West, L. D., Stark, K. E., Murphy, J. J., Lyle, J. M., and F. A. Ochwada-Doyle, 2015 Survey of recreational fishing in New South Wales and the ACT, 2013/14. Fisheries Final Report Series No. 149. NSW Department of Primary Industries, Wollongong.
- Williams, A., Hamer, P., Haddon, M., Robertson, S., Althaus, F., Green, M. and J. Kool, 2016. Determining Blue-eye Trevalla stock structure and improving methods for stock assessment, FRDC project 2013/015, FRDC, Canberra. 124p.
- Williams, A, Hamer, P, Haddon, M, Robertson, S, Althaus, F, Green, M and J, Kool, 2017. Determining blue-eye trevalla stock structure and improving methods for stock assessment, Fisheries Research and Development Corporation, Canberra.
- Wayte, S., Dowdney, J., Williams, A., Bulman, C., Sporcic, M., Fuller, M. and A. Smith, 2007. Ecological Risk Assessment for the Effects of Fishing: Report for the otter trawl sub-fishery of the Commonwealth trawl sector of the Southern and Eastern Scalefish and Shark Fishery. Report for the Australian Fisheries Management Authority, Canberra.

Appendix 1

Reliability and Relevance of the Commonwealth Assessment to assessment of stock status in NSW

The current Commonwealth assessment of Blue-eye Trevalla could adequately inform the decision process for an NSW TAC determination, accepting and understanding the limitations and restrictions of the Commonwealth assessment framework and understanding that data collected and analysed at a finer resolution and considered within the NSW jurisdiction could deliver some outcomes inconsistent with the Commonwealth assessment. The benefits of adopting Commonwealth assessments include the application of processes exposed to broad review, including by management, science and industry representatives within the Commonwealth fishing sector, as well as observers from other stakeholder groups (e.g. NSW DPI Fisheries). The Commonwealth assessments have not, however, been developed to provide specific outputs for jurisdictions other than the Commonwealth and do not necessarily include or apply data at resolutions more applicable to alternate jurisdictions. Therefore, applying these assessments to inform NSW total allowable catch (TAC) determinations is done understanding that there are limitations in the data used and the application of the data to a scale other than that to which the assessment was applied.

Moreover, applying the assessment of Blue-eye Trevalla from the Commonwealth to inform the status of NSW Blue-eye Trevalla and reliably inform management decisions for this species assumes (among other issues) that the assessment represents the same population(s) being harvested by these fisheries. Support for this assumption is provided by the 2020 determination of the stock structure of Blue-eye Trevalla for the 2020 SAFS reports as a management unit at the scale of eastern Australia (Emery et al. 2020).

NSW and Commonwealth SESSF catch rates

Trends in standardised catch rates (CPUE) for Blue-eye Trevalla taken by dropline in the Ocean Trap and Line, Line East fishery operating within NSW jurisdiction (Figure 7) are generally consistent with indices of abundance based on spawning stock biomass trends in Commonwealth Trawl Sector and Scalefish Hook Sector (Figure 2). Both sources of data, show decreasing trends in abundance between 2019 and 2013, with similar increases in abundance peaking in 2015. Abundance subsequently decreased to the lowest point in 2019 (extent of commonwealth CPUE). Further investigation is required to assess Commonwealth SESSF catch rates trends post 2019 to determine whether they are similar to those recent trends in CPUE observed in the NSW OTLLE fishery. The trends in CPUE for Blue-eye Trevalla taken in the OTLLE fishery and within SESSF suggests that the component of the stock in NSW waters is currently exhibiting similar dynamics (with respect to abundance) to the component of stock under Commonwealth jurisdiction and this is consistent with the assumption of a single biological stock.

Appendix 2

NSW catch statistics and additional information relevant to TAC setting in NSW

Information presented in figures and tables below is summarised by fiscal year (July to June). Commercial fishery data presented in this section is limited to data from the Ocean Trap and Line Fishery from 2009/10, as contemporary supplementary information to the assessment and to inform NSW TAC determinations. The exception to this the reported total catches of Blue-eye Trevalla from all fishing methods from 1998/99. Data reporting total catch and catch of different

gear types within the OTLLE endorsement from 2009/10 have been sourced from the NSW DPI database FishOnline.

NSW commercial fishery records have not been consistently reported throughout the history of the fishery. Notably, between 1997/98 and 2008/09 (inclusive), fishers reported monthly catch and effort (in days) for each fishing method (gear type). From 2009/10, monthly reports of daily catch and effort (hours) and fishing method have been required. To construct a longer time series of data (from 1997/98 to present), daily records from 2009/10 are re-aggregated into monthly catches (kg) by fisher and gear type, with effort in days per month estimated from the number of distinct fishing dates in each month where the method was reported and where there was a reported landing of the species of interest in that month, irrespective of whether the species was reported on each of the days, to be consistent with earlier reporting.

State-wide fisheries catch

Annual catches of Blue-eye Trevalla have generally declined over the last two decades, from over 100 t in the late 1990s to < 10 t in the last two years (Figure 3). In 2020/21, the total catch of Blue-eye Trevalla was 7.92 t (Figure 3). Annual catches are dominated, almost exclusively ($\geq 97\%$) by those from the NSW OTLLE endorsement (Figure 4). In the 2020/21 season, total landings of Blue-eye Trevalla in NSW were 3.5% of the total commercial landings from the Commonwealth fisheries (224.6 t; Southern and Eastern Scalefish and Shark Fishery SESSF).

Ocean Trap and Line Fishery catch and catch rate

Within the OTLLE fishery, Blue-eye Trevalla are caught predominantly using dropline, handline and setline fishing methods (Figure 5). Dropline dominates the catch (2009/10-2020/21; range 3.17–38.8 t.yr⁻¹) followed by handline (2009/10-2020/21; 3.17-30.8 t.yr⁻¹). Since 2009/10, setline gear has averaged about 2.6 t.yr⁻¹, equating to about 14% of the total OTLLE catch (Figure 5). Within the general setline gear, setline demersal (STD) is responsible for most of the catch (range 1.49–5.3 t.yr⁻¹), with catch by trotline gear in 2009 recorded as 2.3 t, although this is likely to be associated with misreporting of method from one fisher.

Levels of catch and effort (days), where a consistent effort series is available, have been declining since at least the late 1990s (Figure 6). Prior to 2008/09, annual catches exceed 40 t.yr⁻¹ (range 41–118 t.yr⁻¹) and effort (days) exceed 700 days.yr⁻¹ (range 775–1538 days.yr⁻¹). Since 2009/10, catches have declined from > 30 t.yr⁻¹ to < 5 t.yr⁻¹. In 2020/21 catch and effort (days) was 4.2 t and 46 days, respectively and the lowest recorded levels in the history of the fishery (Figure 7). Commensurate declines in catch and effort through time have resulted in a generally stable time series of CPUE (kg.day⁻¹), although a substantial decline reported between 2012/13 and 2013/14, and low levels of CPUE in recent years suggest stocks may be less stable and/or fleet behaviour and dynamics has changed.

Macbeth and Gray (2015), from fishery-dependent observer days, reported Blue-eye Trevalla comprised the greatest proportion of dropline catch in the NSW OTL Fishery (23.4% of the catch by number) and in the northern region (19.9% of the catch by number), with > 99% of the dropline catch of Blue-eye Trevalla being retained. The catch rate (fish per dropline day) of Blue-eye Trevalla reported by Macbeth and Gray (2015) ranged between 12.5–20.5 fish per dropline day in the south region, and 2.8–7.3 fish per dropline day in the north region. In addition, size-class frequency distributions of Blue-eye Trevalla from observed dropline days fished, indicate substantially larger and fewer fish in the north and central regions of the state (modal size class 75–79 cm FL) compared with smaller but more frequent fish in the south region (50–54 cm FL) (Macbeth and Gray 2015).

Additional information relevant to TAC setting in NSW

1. The NSW Blue-eye Trevalla TAC for the May 2020-April 2021 fishing season was set at the 5-year maximum catch of 30.0 tonnes.
2. The Independent Allocation Panel (IAP) recommended that the initial allocation of quota shares for Blue-eye Trevalla be calculated based on 20% on the proportion of access shares held + 80% on recorded landings for an individual fishing business in the Ocean Trap & Line – Line East Share Class over the selected criteria period 2009/2010 to 2016/2017 (inclusive), but with the “worst catch year” for each business removed (https://www.dpi.nsw.gov.au/data/assets/pdf_file/0009/832464/Ocean-Trap-and-Line-IAP-Final-Report.pdf).
3. Statistics describing landings of Blue-eye Trevalla from NSW commercial fisheries may inform determination of a NSW TAC that is consistent with the development an inter-jurisdictional resource sharing policy.
4. Landings (quota usage) of 7.92 t were reported against a TAC of 30.0 t in 2020/21 which suggests that the current TAC was not constraining total catches.
5. In 2020/21 approximately 11.1 t of quota was held by fishing business that reported nil landings of Blue-eye Trevalla. But note, of this 11 t, 3.6 t was transferred to other fishers. e.g. Fishers with zero allocation caught 2.3 tonnes, by transferring in quota from other FBs.
6. 7.29 t (24.3%) of the 2021/22 Blue-eye Trevalla TAC (30.0 t) was taken in NSW at 1st November 2020 (51% of season complete).
7. SESSF TAC recommendation for Blue-eye Trevalla for 2021/22 was 241 t, which was a 207 t reduction on the 2020-21 TAC (448 t, https://www.afma.gov.au/sites/default/files/sessf_tac_recommendations_2021-22_-_for_concession_holders.pdf).

Stock Status Summary – 2021/22



NSW Stock Status Summary – Blue-eye Trevalla (*Hyperoglyphe antarctica*)

Figure 3. Annual catch (t) of Blue-eye Trevalla from all fishing methods reported to NSW from 1998/89 to 2020/21.





Figure 4. Annual catch (t) of Blue-eye Trevalla from NSW Ocean Trap and Line – Line East (grey; OTLLE) and all other endorsement codes (black, OTHER) from 2009/10 to 2020/21.



Stock Status Summary – 2021/22



NSW Stock Status Summary – Blue-eye Trevalla (*Hyperglyphe antarctica*)

Figure 5. Annual catch of Blue-eye Trevalla from NSW Ocean Trap and Line - Line East (OTLLE) – Dropline (dark grey; DPL), Setline (white; demersal, trotline and unspecified setline), Handline (light grey) and all other methods (black) from 2009/10 to 2020/21.





Figure 6. Dropline only - Annual effort (days) using dropline in the OTLLE from 1998/89 to 2020/21.



Stock Status Summary – 2021/22



NSW Stock Status Summary – Blue-eye Trevalla
(*Hyperoglyphe antarctica*)

Figure 7. Dropline only - Annual standardised CPUE (kg.day⁻¹ from daily records) using dropline in the OTLLE from 2009/10 to 2020/21. The horizontal line represents the average catch rate (2009/10 to 2020/21). Vertical line = TAC introduction.

