

Assessment Authors and Year

Stewart, J. 2023. NSW Stock Status Summary 2022/23 – Blue Mackerel (*Scomber australasicus*). NSW Department of Primary Industries, Fisheries. 12 pp.

Stock Status

Current stock status	On the basis of the evidence contained within this assessment, Blue Mackerel is currently assessed as sustainable .
----------------------	--

Stock structure & distribution

Blue Mackerel occur in sub-tropical and temperate waters of the Pacific and Indian Oceans. They inhabit inshore and continental shelf waters off all states of Australia except the Northern Territory. The stock structure of Blue Mackerel is uncertain (Patterson et al., 2022), with genetic analysis of samples from southern Queensland, Western Australia and New Zealand indicating population subdivision with differences detected between Western Australia and Queensland, and between Western Australia and New Zealand, but not between Queensland and New Zealand (Ward et al., 2007; Whittington et al., 2012). Blue Mackerel off southern Australia is currently considered to be comprised of two biological stocks: the Western stock that extends from western Tasmania to southern Western Australia and the Eastern stock, which occurs to the east of Bass Strait. The Commonwealth Small Pelagic Fishery (SPF) manages and assesses Blue Mackerel along eastern Australia as a single east coast stock.

This stock status summary details stock assessment results and relevant fisheries statistics to inform the setting of a Total Allowable Catch (TAC) for the NSW Ocean Hauling Purse Seine (OHPS) Blue Mackerel quota fishery for the 2023/24 fishing season (May to April). Assessment of stock status for Blue Mackerel is principally based on the Commonwealth SPF derived assessment that utilizes estimates of spawning biomass from periodic egg surveys using the Daily Egg Production Method (DEPM). Given that the DEPM-based assessment is for the entire eastern stock, the results are appropriate for application to the NSW component of the stock.

Biology


Blue Mackerel grow relatively quickly, reaching approximately 17-22 cm FL after one year, and have been recorded to attain a maximum age of about 8 years and length of 44 cm FL in Australian waters. Blue Mackerel have been aged to be more than 20 years old in New Zealand. Blue Mackerel mature at about 24-28 cm fork length (FL) and spawning is thought to occur during late winter and spring in outer continental shelf waters off northern NSW and southern QLD. Eggs and larvae have been observed in high abundances in shelf waters with temperatures of 18 to 21°C (Neira and Keane 2008). Reproductive parameters for Blue Mackerel in Eastern Australia are not well understood.

FISHERY STATISTICS

Catch information

Commercial

Most of the eastern Blue Mackerel catch was historically taken in state fisheries; however since 2015/16 the Commonwealth catch has exceeded state commercial catches. The total combined commercial NSW and Commonwealth catch in 2020/21 increased to 6,662 t, comprising 6,215 t from the Commonwealth and 447 t from NSW state fisheries. In 2021/22 the Commonwealth catch increased substantially to 10,051 t (Fig. 1), and the NSW reported catch at the time data was extracted was only 134 t, (Patterson et al., 2022). Landings from NSW estimated from quota usage (rather than official logbook data) during 2021/22 were likely approaching 500 t (comprising approximately 291 t from ocean hauling, 36 t from Commonwealth tuna boats under permit and recreational catch) (Fig. 2). Total harvests during 2021/22 were therefore likely approaching 10,500 t and the highest on record.



Note: TAC Total allowable catch.

Figure 1 Commonwealth eastern Blue Mackerel catch and TAC 2003/04 fishing season to 2021/22 fishing season (from Patterson et al. 2022).

Stock Status Summary – 2022/23



NSW Stock Status Summary – Blue Mackerel (*Scomber australasicus*)




Figure 2. Reported landings of Blue Mackerel by the NSW commercial fishery 1955/56 to 2021/22. Data excludes catch reported under section 37 permit for Commonwealth tuna bait.

Information on commercial discarding of Blue Mackerel is limited. Blue Mackerel are rarely released in the commercial line fishery (MacBeth and Gray 2013). Survival of released fish from purse-seine operations is thought to be high (industry pers. comm).

Recreational & Charter boat

Blue Mackerel are very important to recreational and charter fishers, and the recreational harvest is significant but poorly estimated. The most recent surveys of recreational harvest in NSW by 1 and 3 year licence holders and housemates were done during 2017/18 and 2019/20. These estimated the Blue Mackerel harvest at approximately 41 t and 51 t respectively (Murphy et al., 2020, NSW DPI Unpublished data). Approximated conversions to make these estimates comparable to the 2000/01 state-wide survey that encompassed total recreational harvest scales these to be around 125 t and 156 t and approximately 21% and 23% of the total harvest in NSW waters.

Approximately 20% of Blue Mackerel in recreational catches were estimated to have been released during 2017/18 and during 2019/20, with unknown survival (Murphy et al., 2020, Murphy et al., 2022).

Indigenous

There is no information available on the Aboriginal catch of Blue Mackerel in NSW waters.

Illegal, Unregulated and Unreported

The level of Illegal Unregulated and Unreported (IUU) fishing is unknown.

Fishing effort information

Days fished in the NSW purse seine fishery when Blue Mackerel were landed remained relatively stable between 2009/10 and 2017/18, fluctuating between approximately 400 and 600 days per year. Days of effort increased to be between around 650 to 850 between 2017/18 and 2019/20, but declined to approximately 500 during 2020/21 and then to only 249 during 2021/22. (Fig. 3).




Figure 3. Effort (days fished) for purse seine fishers that reported landing Blue Mackerel in NSW 2009/10 to 2021/22.

Catch Rate information

Catch rates of Blue Mackerel (median kg per day purse seining in the Ocean Hauling Fishery) have fluctuated since 2009/10 with higher rates during 2019/20 and 2020/21, before falling dramatically to approximately 160 kg/day during 2021/22 (Fig. 4). This dramatic decline appears to have been driven partly by regular reporting of small quantities of Blue Mackerel, presumably taken as byproduct, and non-reporting by some of the larger quota holders in the fishery during 2020/21. More detailed logbook data in terms of catch per shot indicates considerable annual increases since 2013/14, with data for 2021/22 showing the highest catch rate per shot (Fig. 5), noting these are raw catch rates and do not account for factors such as vessel size, net size etc. At present such catch rate standardizations are not required for assessment purposes. It should be noted that catch rates of Blue Mackerel by the method of purse seine are unlikely to be useful for inferring relative abundance due to: (i) the schooling nature of the species; (ii) the ability of the gear to encircle entire schools of fish, and; (iii) the market driven nature of fishing operations for this species in NSW.

Stock Status Summary – 2022/23



NSW Stock Status Summary – Blue Mackerel (*Scomber australasicus*)




Figure 4. Commercial catch rates (raw kgs per day) of Blue Mackerel using purse seining for years 1997/98 to 2021/22 in NSW.




Figure 5. Commercial catch rates (kgs per shot) of Blue Mackerel using purse seining for years 2009/10 to 2021/22 in NSW.

STOCK ASSESSMENT

Blue Mackerel is assessed in terms of biomass through Daily Egg Production Method (DEPM) surveys. Sustainable levels of fishing are assessed through catch as a percentage of biomass. Management Strategy Evaluation (MSE) of the Commonwealth SPF Harvest Strategy (Smith et al., 2015) established that an exploitation rate of up to 23% may be suitable for eastern Blue Mackerel under the Commonwealth SPF harvest strategy. The current harvest strategy exploitation rate following a Tier 1 assessment is only 15%, applied to maintain the stock in the vicinity of the target reference point of 50% of unfished levels and therefore ensuring a very low probability of the stock falling below 20% of unfished levels. A Stochastic Stock Reduction Analysis (SSRA) was also done to estimate depletion levels up to 2015 (Punt et al., 2016a; Punt et al., 2016b).

Stock Assessment Methodology

Year of most recent assessment:

2022

2019 - Daily Egg Production Method (DEPM) biomass estimate.

2015 - Stochastic Stock Reduction Analysis (SSRA).

Assessment method:

Daily Egg Production Method (DEPM) biomass estimate (Ward et al., 2021).

Stochastic Stock Reduction Analysis (SSRA) and Management Strategy Evaluation (MSE) of the Commonwealth SPF Harvest Strategy (Smith et al., 2016; Punt et al., 2016a; Punt et al., 2016b).

Main data inputs:

Egg survey during September 2019 between Sandy Cape, Queensland and Ulladulla, NSW. The survey produced estimates of Blue Mackerel egg abundance, egg age and spawning area.

Adult reproductive parameters: average weight, sex ratio, batch fecundity, spawning fraction.

Catch and effort data.

SSRA: Catch, 2014 spawning biomass estimate, growth, maturity, selectivity, stock-recruitment relationship.

MSE: Weight, maturity and selectivity by age.

Key model structure & assumptions:

SSRA: age-structured model, fixed parameters for weight-at-age, natural mortality, selectivity at age and stock-recruitment steepness. Free parameters unfished recruitment, fishing mortality on fully-selected age classes, deviations around the stock-recruitment relationship. 2014 spawning biomass estimate based on the DEPM derived 83,300 t with a CV of 0.5. Assumptions include negligible catch prior to 1997/98, and that assumed parameters are correct.

MSE operating model is age-structured, and recruitment is driven by spawning stock biomass and uses pre-specified values for biological parameters (natural mortality, growth, maturity, and stock-recruit steepness).

Stock Status Summary – 2022/23



NSW Stock Status Summary – Blue Mackerel (*Scomber australasicus*)

Sources of uncertainty evaluated:

Considerable uncertainty exists around all of the key input data for the Blue Mackerel DEPM assessment. Sensitivity analyses were done for all parameters to determine which had the largest influence on estimated spawning biomass. These were done by varying each individual parameter whilst keeping the others constant at the value used to calculate spawning biomass.

Conclusions were drawn based on the most precautionary parameter estimates, resulting in the spawning biomass likely to be under-estimated.

MSE testing of the SPF harvest strategy rules to examine the probability of the biomass falling below the limit reference point of 20% of unfished levels with a less than 10% chance over 50 years.

Status Indicators - Limit & Target Reference Levels

Biomass indicator or proxy	DEPM derived estimate of spawning biomass.
Biomass Limit Reference Point	20% of unfished levels with a less than 10% chance.
Biomass Target Reference Point	50% of unfished levels
Fishing mortality indicator or proxy	Catch as a proportion of spawning biomass.
Fishing mortality Limit Reference Point	<p>Annual catch is less than 15% of the DEPM derived estimate of spawning biomass. This is the Tier 1 exploitation rate in the Commonwealth SPF Harvest Strategy for setting a Recommended Biological Catch (RBC) for each of five fishing seasons following a DEPM assessment.</p> <p>Five years after a Tier 1 assessment, the RBC is set at the Tier 2 level that is 7.5% of the DEPM derived estimate of spawning biomass.</p> <p>Five years after a Tier 2 assessment, if no updated DEPM is done, the RBC is set at the Tier 3 level that is 3.75% of the DEPM derived estimate of spawning biomass.</p>
Fishing Mortality Target Reference Point	As above. The exploitation rate is applied to maintain the stock around the target reference level of 50% of unfished biomass.

Stock Assessment Results

Recent harvests of east coast Blue Mackerel have been increasing, but remain below the reference level of 15% of the 2019 derived DEPM estimate of spawning biomass (rounded down to 80,000 t for calculation of the RBC in recognition of the imprecision inherent in such surveys)

Stock Status Summary – 2022/23



NSW Stock Status Summary – Blue Mackerel (*Scomber australasicus*)

(Ward et al., 2021) with the RBC calculated as 15% x 80,000 t ~12,000 t (Patterson et al., 2022). This level of fishing mortality is unlikely to have substantially reduced the spawning biomass.

Stock Assessment Result Summary

Biomass status in relation to Limit	Stochastic Stock Reduction Analysis (Punt et al., 2016a; Punt et al., 2016b) estimated that the 2015 depletion of Blue Mackerel was likely to be fairly close to the average unfished level. The SPF Harvest Strategy exploitation rates have been tested to provide a high likelihood that stocks will be maintained, on average, at the target reference point of 50 per cent of unfished levels, with a less than a 10 per cent chance over 50 years of falling below the limit reference point of 20 per cent of unfished levels.
Biomass status in relation to Target	As above
Fishing mortality in relation to Limit	Recent harvests of east coast Blue Mackerel have been increasing, but remain below the reference level of 15% of the 2019 derived DEPM estimate of spawning biomass (rounded down to 80,000 t for calculation of the RBC in recognition of the imprecision inherent in such surveys) (Ward et al., 2021) with the RBC calculated as 15% x 80,000 t ~12,000 t.
Fishing mortality in relation to Target	As above
Current SAFS stock status	Sustainable in 2020
Current Commonwealth stock status	Not overfished and not subject to overfishing

Fishery interactions

Commonwealth Small Pelagic Fishery – purse seine and midwater trawl, interacts with the NSW commercial fishery.

Commonwealth Tuna boats accessing Blue Mackerel for bait under permit.

Recreational fishers who harvest Blue Mackerel for food and bait. An historically contentious fishery interaction with concerns about the impact of commercial operations on the availability of bait for recreational fishers and on the distribution and therefore availability of gamefish.

Qualifying Comments

The DEPM-based estimates of Blue Mackerel spawning biomass are highly likely to be under-estimates, due to any potential biases in terms of key parameters (such as spawning area and the assumption that surveys are done at the peak spawning time) always leading to under-estimating

spawning biomass. A recognised uncertainty in the Blue Mackerel DEPM biomass estimation has been the use of adult reproductive parameters that were obtained from fish collected in South Australia during the early 2000s. AFMA are currently funding a pilot study to investigate the presence of Blue Mackerel in spawning condition off the east-coast.

The very wide confidence intervals of DEPM-derived spawning biomass need to be acknowledged. In 2019 it was estimated to be 88,265 t (95% confidence intervals 33,320 to 143,209 t) Ward et al., 2021. In 2014 it was estimated at 83 300 t with 95% confidence intervals between 35,100 and 165,000 t (Ward et al., 2015), and in 2004 it was estimated at 23,009 t with 95% confidence intervals between 7,565 and 116,395 t (Ward and Rogers, 2007).

It should be noted that a component of the Stochastic Stock Reduction Analysis (Punt et al., 2016a) investigated the effect of temporal auto-correlation in recruitment (alternating periods of high or low recruitment and something that is observed in other small pelagic species) and concluded that “it is not possible to maintain stocks above the reference points considered with the pre-specified probability (10%) even without fishing, but this is not the case if deviations in recruitment about the stock-recruitment relationship are temporally uncorrelated.” Such a finding supports the rationale of implementing precautionary harvest strategies that rely on regular biomass estimates.

References

- Macbeth, W.G. and Gray, C.A., 2016. Observer-based study of commercial line fishing in waters off New South Wales. Fisheries Final Report Series, (148,151).
- Murphy, J.J., Ochwada-Doyle, F.A., West, L.D., Stark, K.E. and Hughes, J.M., 2020, The NSW Recreational Fisheries Monitoring Program - survey of recreational fishing, 2017/18. NSW DPI - Fisheries Final Report Series No. 158.
- Murphy, J. J., Ochwada-Doyle, F. A., West, L. D., Stark, K. E., Hughes, J. M., Taylor, M.D., 2022. Survey of recreational fishing in NSW, 2019/20 – Key Results. NSW DPI – Fisheries Final Report Series No. 161. ISSN 2204-8669
- Patterson, H, Bromhead, D, Galeano, D, Larcombe, J, Timmiss, T, Woodhams, J and Curtotti, R 2022, Fishery status reports 2022, Australian Bureau of Agricultural and Resource Economics and Sciences, Canberra. CC BY 4.0. <https://doi.org/10.25814/gx9r-3n90>.
- Punt, A.E., Little, R. and Hillary, R. 2016a. Updated management strategy evaluation for Eastern Jack and Blue Mackerel, report for AFMA, CSIRO Oceans and Atmosphere Flagship, Hobart.
- Punt, A.E., Little, L.R. and R. Hillary. 2016b. Assessment for Eastern Blue Mackerel. Document submitted to AFMA. 6pp.
- Smith, A., Ward, T., Hurtado, F., Klaer, N., Fulton, E. and Punt, A. 2015. Review and update of harvest strategy settings for the Commonwealth Small Pelagic Fishery: single species and ecosystem considerations, report for FRDC project 2013/028, CSIRO Oceans and Atmosphere Flagship, Hobart.
- Ward, T. M., Grammar, G. L. and Ivey, A. R. (2021). Spawning biomass of Blue Mackerel (*Scomber australasicus*) and Australian Sardine (*Sardinops sagax*) in the East sub-area of the Small Pelagic Fishery. Report to the Australian Fisheries Management Authority. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2021/000047-1. SARDI Research Report Series No 1087. 54pp.

Ward, T.M., Grammer, G., Ivey, A., Carroll, J., Keane, J., Stewart, J. and Litherland, L. 2015. Egg distribution, reproductive parameters and spawning biomass of blue mackerel, Australian sardine and tailor off the east coast during late winter and early spring, FRDC project 2014/033, FRDC & SARDI, West Beach.

Ward, T.M. and Rogers, P.J. 2007. Evaluating the application of egg-based stock assessment methods for blue mackerel, *Scomber australasicus*, in southern Australia, Final report to FRDC, Canberra.

Appendix 1

Blue Mackerel catch reporting by the NSW Ocean Hauling Purse Seine fishery.

Currently NSW fishers landing Blue Mackerel under quota are required to report their landings through two separate processes. Fishers are required to report 'real time' quota use through the FisherMobile application. In addition, fishers are required to submit details of their fishing activity, including catch and effort, through mandated logbooks (paper-based or via the FisherDirect online reporting portal). Monitoring and assessment is done based on the time series of reported catch and effort information submitted by fishers through their mandated logbooks.

Since the introduction of quota management for the 2019/20 fishing season (May to April) the quantity of Blue Mackerel reported as being landed through compulsory logbooks has been less than the quantity reported through the quota management system (Fig. 6). During 2021/22 57% of the reported quota used was reported through the logbooks. It is important to note that the catch and effort data used for assessments, and supplied to the Commonwealth for use in TAC setting for their fisheries, is derived from the mandatory NSW logbooks, and not from the real-time quota reports.




Figure 6. Blue Mackerel reported as landed through mandatory logbooks and real-time quota reporting since the introduction of quota management. Note that the 2022/23 quota season was only 8.5 months completed at the time of data extraction, and that logbook returns were likely yet to be compiled.

Performance of the NSW Blue Mackerel quota fishery against the allocated TAC of approximately 758 t p.a. is relatively low, whether assessed from real-time quota reports (Fig. 7) or mandatory logbooks (Fig. 8). The official catch data indicates that less than 20% of the allocated quota for 2021/22 was caught, yet quota reports indicate that this is a large under-estimation.

Stock Status Summary – 2022/23



NSW Stock Status Summary – Blue Mackerel (*Scomber australasicus*)




Figure 7. Blue Mackerel landings reported through the real time quota usage system, and allocated TAC, since quota management was implemented. Note that the 2022/23 fishing season was only 8.5 months completed when these data were collated.




Figure 8. Blue Mackerel landings (financial year) reported through the mandatory logbooks, and allocated TAC, each since 2009/10. Note data for 2022/23 is incomplete. Quota management began in 2019/20.