

Sheep health - scabby mouth

September 2016, Primefact A3.9.41, third edition

J.W Plant Former Special Veterinary Officer Sheep Health
Geoff Casburn Sheep Development Officer, Wagga Wagga

Scabby mouth is a highly contagious viral disease of the skin of sheep and goats which usually affects lambs and kids in their first year of life.

Infection usually causes scabs and pustules around the lips (figure 1), but it can also affect the udder and the skin around the coronet and pasterns.

Once the disease is introduced to a property, the virus will remain in the environment for many years, but may not cause infection every year.

Scabby mouth can be prevented by vaccination.

Figure 1 Lamb affected with scabby mouth showing lesions around the lips and nostrils.



The disease is caused by a pox virus which is normally present in scabs on affected animals. The virus can survive off the sheep for many years under the right conditions.

Clinical signs

Sheep become infected through injuries or abrasions on the skin of the muzzle, the lips or coronet or other areas of bare skin.

Two to four days after infection the skin becomes reddened and may show some swelling. The first signs of infection are usually seen near the corners of the lips. The next stage is the

development of blisters which quickly develop into pustules. These pustules then rupture, excreting virus and forming scabs, giving the disease its typical appearance.

The scabs lift in one to two weeks, leaving raw skin which quickly heals and returns to normal. In some lambs, the lesions spread along the lips and merge to form scabs which may involve the whole of the lips. In severe cases, the lesions may extend into the mouth, onto the tongue and into the nostrils.

The lesions usually disappear within four to six weeks of the initial infection.

Severely affected lambs may be unable to feed for several days, resulting in a loss of condition and an increased susceptibility to other diseases or death by starvation in young lambs. **Scabby mouth alone does not cause deaths, but it will severely depress growth rates.**

Lesions on the teats and udders of ewes may predispose to mastitis. These lesions are painful and the ewe may not allow the lamb to drink.

Lesions can also develop on the scrotum of rams, causing a thickening of the scrotal skin, affecting semen quality by interfering with the cooling mechanism of the testes.

Scabby mouth infection can also occur around the coronet and pasterns, potentially leading to secondary infection such as 'strawberry footrot' and severe lameness.

Transmission of infection

The virus will survive for months in exposed areas and can survive for years in a dry sheltered area such as around sheds or in yards. The virus, although present, will not cause any problems until it gains entry to injuries on the skin of susceptible animals.

The common cause of injury is grazing in paddocks with a lot of thistle or burr or handfeeding hay containing thistle or burr. Grass seeds and stubble can also cause the initial injury.

Outbreaks in housed sheep can be more severe because of the congregation of animals in a confined space and the restricted feeding areas.

Contamination of feed troughs will result in a rapid spread of infection through a pen of susceptible sheep, particularly if they are being fed on hay containing some thistle.

Scabby mouth infection of the foot usually develops when sheep are running in wet lush pasture containing thistles. These conditions predispose the feet, coronet and pasterns to injury, making them susceptible to infection.

Diagnosis

The disease is best diagnosed by a clinical examination of affected animals. In some animals, both dermatophilosis and scabby mouth may occur together. Conditions which can be confused with scabby mouth include dermatophilosis (figure 2), bacterial skin infections and photosensitisation.

Figure 2 Lamb affected with dermatophilosis around nose. Note absence of lesions around lips and nostrils.



No treatment will kill the virus and the disease normally runs its course in three to four weeks. Spread may be reduced by moving affected mobs to paddocks where injury to the skin is less likely, such as paddocks with less thistle or burr. Severely affected animals may need treatment to overcome secondary infections.

Control

Sheep which recover from scabby mouth have a lifelong immunity against severe infection. However, they are susceptible to reinfection, but the infection is milder and of shorter duration and may go unnoticed.

Sheep are best vaccinated as lambs at lamb marking. This involves scratching the live vaccine onto the skin on the inner thigh. The lamb will develop a skin infection at the site of vaccination within five to ten days and develop immunity in 14 days.

Sheep should be checked five to ten days after vaccination to ensure that the vaccine has taken. Successful vaccination produces pustules along the line of the scratches. These pustules progress to small scabs which normally disappear three to four weeks after vaccination.

Pregnant ewes should be vaccinated behind the elbow. If ewes are vaccinated on the inner thigh, then the udder may become infected resulting in rapid transmission of infection to highly susceptible young lambs.

Vaccination should not be considered on properties unless scabby mouth is causing problems. If sheep on an uninfected property are vaccinated, then the virus is introduced on to the property and a regular vaccination program may have to be carried out to protect stock in future years.

Handling of the Vaccine

The vaccine contains a live virus and must be handled carefully to maintain its effectiveness. It should be stored in the fridge (not frozen) between 2° and 8° and in a cold storage bag during transport and use. The vaccine must be kept protected from sunlight.

Care needs to be taken to ensure the vaccine site is not contaminated with antiseptic used during the lamb marking operation.

Handle the vaccine with care as humans can be infected through skin cuts or abrasions. Lesions develop as red swollen nodules on the skin four to six days after infection and usually take a long time to heal

For updates go to
www.dpi.nsw.gov.au/factsheets

© State of New South Wales through the Department of Industry, Skills and Regional Development, 2015. You may copy, distribute and otherwise freely deal with this publication for any purpose, provided that you attribute the NSW Department of Primary Industries as the owner.

Disclaimer: The information contained in this publication is based on knowledge and understanding at the time of writing (September 2016). However, because of advances in knowledge, users are reminded of the need to ensure that information upon which they rely is up to date and to check currency of the information with the appropriate officer of the Department of Primary Industries or the user's independent advisor.

ISSN 1832 6668

INT16/109784