

Blackleg in cattle

Dr Sarah Robson

Regional Animal Health Leader, Animal & Plant Biosecurity, Wagga Wagga

J.M. Wilson

Former District Veterinarian, Rural Lands Protection Board

Introduction

Blackleg is a fatal disease of young cattle. It produces an acute local infection, and the resulting blood poisoning leads to rapid death.

The name 'blackleg' derives from the fact that the site of infection is often a leg muscle, and that the affected muscle is dark in colour.

Occurrence of blackleg

Although the disease is widely distributed in New South Wales, most losses from blackleg occur in the eastern half of the state. In some areas the disease may appear on several properties, while in other areas only isolated farms are affected. Sometimes, only part of a property is affected.

Susceptible animals

Although blackleg has been found in cattle as young as 2 months old, most losses occur in cattle between 6 months and 2 years of age. Occasionally, losses are seen in adult cattle.

Generally, the best conditioned animals are affected, with most losses occurring where there is an abundance of feed.

Blackleg can occur at any time of the year, though more losses are seen during hot, humid weather or following the sudden onset of cold periods.

Cause

Blackleg is produced by spore-forming bacteria. The organisms most commonly responsible are *Clostridium chauvoei* and, less frequently,

C. septicum. Spores produced by the clostridia can lie dormant in the soil for years without losing their potency.

Method of infection

Bacterial spores are eaten in contaminated feed or soil. The spores then enter the bloodstream and lodge in various organs and tissues, including muscles. Here they lie dormant until stimulated to multiply, possibly by some slight injury to the animal. The injury reduces blood flow to the area, thereby reducing the supply of oxygen to the tissues. In the absence of oxygen, the spores germinate and multiply. As they grow, the bacteria produce toxins which destroy surrounding tissues. The toxins are absorbed into the animal's bloodstream which makes the animal acutely sick and causes rapid death.

Signs of the disease

Blackleg should be suspected if an animal aged between 6 months and 2 years:

- becomes lame with swelling of a muscle;
- stops grazing;
- appears sick and quickly goes down.

However, these signs are usually of such short duration that they may be missed. More frequently, a thrifty calf or yearling is simply found dead. Gas is detectable under the skin and this produces a crackling sensation when the skin is rubbed with the hand.

The rapid accumulation of gas under the skin and in the body cavity gives the carcass a bloated appearance, with the limbs spread apart and pointing upwards. There may be a frothy, blood-stained discharge from the mouth, nostrils and anus.

If the skin over the affected area is removed, excess bubbly bloodstained fluid can be seen, and the muscle immediately below will be dark in colour. However, when the affected muscle is inside the carcass, such as when the heart muscle



is affected, no external evidence of the disease is found.

Figure 1 shows diseased skeletal muscle from a heifer that died suddenly from blackleg. The muscle tissue is dark red and has a dry appearance due to gas formation.

Figure 2 shows diseased heart muscle from a 12-month-old heifer that died suddenly from blackleg. Notice the haemorrhage and the 'half cooked' appearance of the diseased muscle tissue.

Because the disease is accompanied by rapid decomposition, the post-mortem changes may be masked by gas formation, especially if examination of the carcass is delayed for more than a few hours.




Figure 1. Diseased skeletal muscle.



Figure 2. Diseased heart muscle
Photos: Roger Cook, NSW DPI

The only effective means of controlling blackleg is by vaccination. Several makes of multivalent vaccine ('5 in 1' or '7 in 1') are available commercially and care should be taken to follow the manufacturer's instructions.

Always read the label

Users of agricultural or veterinary chemical products must always read the label and any Permit before using the product, and strictly comply with the directions on the label and the conditions of any Permit. Users are not absolved from compliance with the directions on the label or the conditions of the Permit by reason of any statement made or not made in this publication.

- Calves should receive two doses of blackleg vaccine. Two vaccinations 1 month apart are essential to provide the best protection.
- A booster vaccination 12 months later should provide lifelong immunity to blackleg.
- It is desirable to give the initial two doses of vaccine before young cattle reach their most susceptible age of six months.
- To await the occurrence of blackleg before vaccinating is unwise, as vaccines take 10–14 days before they begin to provide immunity.

If store cattle are purchased in blackleg areas, it is wise to vaccinate all newly bought young cattle.

Vaccination techniques

The vaccine should be delivered just under the skin – not into the muscle. Draw up a pinch of skin and insert the needle between the skin and the muscle. The loose skin of the neck is convenient for this. (See Primefact 431 *Beef cattle vaccines* for correct vaccination technique.)

Do not save unused parts of bottles or containers of vaccines for future use, as they can become contaminated with undesirable organisms and/or lose their potency. Destroy any vaccine not used within 24 hours of opening.

Vaccination breakdowns

Modern vaccines are produced under conditions of strict quality control by reputable manufacturers. Occasionally, reports are received of apparent failure of vaccines. When investigated, most of these alleged vaccination failures are due to:

- incorrect dosing;
- faulty technique;
- using time-expired vaccine;
- using material from partly used containers;
- vaccine having been subjected to high temperatures during storage or transportation.

Abscesses or large swellings at the site of vaccination are usually the result of:

- lack of hygiene;
- injecting into muscle rather than beneath the skin.

If all these factors can be ruled out, and breakdown is suspected, seek veterinary advice.

Treatment

The speed with which blackleg kills usually makes individual treatment useless. In some cases, however, animals treated early with antibiotics may survive, although they often suffer permanent

deformity due to partial or complete destruction of muscles.

Carcass disposal

Carcasses of animals known to have died from blackleg should not be opened. Opening the carcass can liberate bacteria which will form spores that will contaminate the ground and subsequently infect other cattle.

Also, do not drag carcasses along the ground. If possible, burn or deeply bury the carcasses where they lie.

Remember

- Blackleg is almost entirely preventable by vaccination.
- Read – and heed – the directions on the vaccine labels.
- If in doubt about the cause of deaths in cattle, consult your veterinary practitioner, district veterinarian or veterinary officer.

© State of New South Wales
through NSW Department of Primary Industries 2007

ISSN 1832-6668

Replaces Agfact A0.9.22

Check for updates of this Primefact at:
www.dpi.nsw.gov.au/primefacts

Disclaimer: The information contained in this publication is based on knowledge and understanding at the time of writing (February 2007). However, because of advances in knowledge, users are reminded of the need to ensure that information upon which they rely is up to date and to check currency of the information with the appropriate officer of New South Wales Department of Primary Industries or the user's independent adviser.

Job number 7325