primefact

IRM1 mandarin

March 2020, Primefact 1725, First edition

Dave Monks and Graeme Sanderson, Research Horticulturists, Dareton



Figure 1. An IRM1 mandarin tree.



Figure 2. IRM1 mandarins.

Estimated maturity period

Region	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Riverina												
Sunraysia												

Origin

IRM1 is a low-seeded Murcott mandarin developed by irradiation breeding in Queensland, Australia. It has Plant Breeder's Rights (PBR) protection.

Fruit quality

Table 1. IRM1 mandarin fruit quality* characteristics.

Skin	Yellow to orange. The rind can adhere tightly to the flesh. Skin removes relatively easily once initial break in the skin has begun. Predominantly smooth with some ridging at the stem end of the fruit. Skin is sensitive to sunburn and wind scar.				
Average rind thickness (mm)	1.9				
Internal quality	Rich, sweet flavour with high 'Brix and juice content.				
Average number of seeds	2.4				
Juice per cent (%)	51				
°Brix	14.9				
Acid per cent (%)	1.1				
Brix:acid ratio	13.5				
Average fruit weight (g)	104				
Average fruit diameter (mm)	61				

*Juice quality levels considered adequate for harvest and developed by sequential analysis of fruit from topworked evaluation trees.

Comments

- IRM1 is a sweet fruit with high juice content. Initial fruit quality testing suggests IRM1 is slightly later maturing than IRM2.
- In 2009, the average fruit yield from trees top-worked in 2005 to mature Valencia orange with a citrange rootstock was 31 kg per tree. Hand thinning to balance the crop load accounted for 30% of the fruit produced on the tree, and of this, one third was removed due to sun damage. Average fruit size was 61 mm in diameter as determined from fruit used for quality testing. Fruit grading showed that 63% of the fruit harvested was in the 58–64 mm size range.
- Murcott mandarin types do not seem to reach large fruit sizes in southern Australian climatic conditions. The adoption of IRM1 low-seeded Murcott in Australia has been predominantly by Queensland citrus growers.

Table 2. Average yield per tree* on nursery propagated field trees (Sunraysia).

Dootstook	Average yield per tree (kg)						
Rootstock	2010 (4-y-old trees)	2011 (5-y-old trees)	2012 (6-y-old trees)	2013 (7-y-old trees)			
C35 Citrange	5	51	1	34			
Citrange	33	35	41	10			
Cleopatra	19	15	42	2			
Swingle	39	57	37	57			
Trifoliata	18	42	40	51			
Volkameriana	29	27	31	12			

Table 3. Average yield per tree* on trees top-worked to Valencia orange.

Rootstock	Average yield per tree (kg)						
	2010	2011	2012	2013			
Citrange	64	52	48	44			
Cleopatra	31	56	19	74			
Trifoliata	80	75	56	86			

^{*}Average yield per tree results are from a small number of evaluation trees and should only be used as a general indication of the variety's potential yield.

There has been enquiry from southern Australia growers about IRMI due to its late maturity, larger fruit size and low seed content.

Acknowledgements

Citrus Australia Ltd (CAL)

Department of Primary Industries and Regional Development, WA

Hort Innovation Australia

Queensland Department of Agriculture and Fisheries



This project has been funded by Hort Innovation using the citrus research and development levy and funds from the Australian Government. For more information on the fund and strategic levy investment visit horticulture.com.au

Reference number: PUB20/131

State of New South Wales through the Department of Planning, Industry and Environment ("Department") 2020. The content has been developed by the Department using funds provided by Horticulture Innovation Australia Limited ("Hort Innovation").

The information contained in this publication is based on knowledge and understanding at the time of writing (December 2019) and was generated from field and nursery trees at Dareton Primary Industry Institute, Sunraysia, NSW, unless otherwise stated. Where quantitative data are presented (e.g. % Juice or rind thickness) they are based on measured properties. Where qualitative data are presented (e.g. thorniness or tendency to split), they are based on observations or brief notes recorded in the field.

Because of advances in knowledge, users are reminded of the need to ensure the information upon which they rely is up to date and to check the currency of the information with the appropriate officer of the Department and the user's independent advisor. Any reliance on the contents of the publication (or any part thereof) will be entirely at the user's own risk and neither Hort Innovation nor the Department will be responsible or liable for any loss, damage, cost or expense allegedly arising from any use or non-use of this publication.

Whilst care has been taken in the preparation of this publication, Hort Innovation and the Department make no representations and (to the extent permitted by law) expressly exclude all warranties regarding the accuracy, completeness or currency of the information, recommendations and opinions contained in this publication.